

# Epidemiology of *Chlamydia trachomatis* in the Middle East and north Africa: a systematic review, meta-analysis, and meta-regression

Alex Smolak\*, Hiam Chemaitelly\*, Joumana G Hermez, Nicola Low, Laith J Abu-Raddad



## Summary

**Background** The epidemiology of *Chlamydia trachomatis* in the Middle East and north Africa is poorly understood. We aimed to provide a comprehensive epidemiological assessment of *C trachomatis* infection in the Middle East and north Africa.

**Methods** We did a systematic review of *C trachomatis* infection as well as a meta-analysis and meta-regression of *C trachomatis* prevalence. We searched PubMed and Embase, as well as regional and national databases up to March 13, 2019, using broad search terms with no language or year restrictions. Any document or report including biological measures for *C trachomatis* prevalence or incidence was eligible for inclusion. We extracted all measures of current (genital or rectal), recent, and ever infection with *C trachomatis*. We estimated pooled average prevalence in different populations using random-effects meta-analysis. Factors associated with prevalence and sources of between-study heterogeneity were determined using meta-regression.

**Findings** We identified a total of 1531 citations, of which 255 reports contributed to 552 *C trachomatis* prevalence measures from 20 countries. No incidence measures were identified. Pooled prevalence of current genital infection was 3·0% (95% CI 2·3–3·8) in general populations, 2·8% (1·0–5·2) in intermediate-risk populations, 13·2% (7·2–20·7) in female sex workers, 11·3% (9·0–13·7) in infertility clinic attendees, 12·4% (7·9–17·7) in women with miscarriage, 12·4% (9·4–15·7) in symptomatic women, and 17·4% (12·5–22·8) in symptomatic men. Pooled prevalence of current rectal infection was 7·7% (4·2–12·0) in men who have sex with men. Substantial between-study heterogeneity was found. Multivariable meta-regression explained 29·0% of variation. Population type was most strongly associated with prevalence. Additional associations were found with assay type, sample size, country, and sex, but not with sampling methodology or response rate (about 90% of studies used convenience sampling and >75% had unclear response rate). There was no evidence for temporal variation in prevalence between 1982 and 2018.

**Interpretation** *C trachomatis* prevalence in the Middle East and north Africa is similar to other regions, but higher than expected given its sexually conservative norms. High prevalence in infertility clinic attendees and in women with miscarriage suggests a potential role for *C trachomatis* in poor reproductive health outcomes in this region.

**Funding** National Priorities Research Program from the Qatar National Research Fund (a member of Qatar Foundation).

**Copyright** © 2019 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

## Introduction

With more than 100 million incident infections every year,<sup>1</sup> *Chlamydia trachomatis* is one of the most common sexually transmitted infections (STIs) worldwide.<sup>2,3</sup> Although curable, control and early detection of *C trachomatis* infection are challenged by its largely asymptomatic nature.<sup>4</sup> Untreated *C trachomatis* infection is associated with serious reproductive tract conditions including pelvic inflammatory disease, ectopic pregnancy, infertility among women, and epididymitis among men.<sup>5,6</sup>

Despite burdensome sequelae, STI control has long languished on health policy agendas. The 2030 Agenda on Sustainable Development<sup>7</sup> aims to remedy this situation and led to WHO's Global Health Sector Strategy on STIs.<sup>8</sup> The strategy proposes an integrated approach for STI prevention and control that addresses core Sustainable Development Goals, mainly through securing

universal access to sexual and reproductive health-care services and rights.<sup>7,8</sup> The first strategic direction of this STI Strategy is “the need to understand the sexually transmitted infection epidemic and response as a basis for advocacy, political commitment, national planning, resource mobilization and allocation, implementation, and programme improvement.”<sup>8</sup>

The epidemiology of STIs, including *C trachomatis*, remains poorly understood in the Middle East and north Africa—a region comprising 10% of the world's population.<sup>9–11</sup> Here, political and sociocultural sensitivities have set STIs low on countries' public health agendas, resulting in limited capacity for surveillance and programmes targeting sexual health, despite the possibility of a hidden disease burden.<sup>9</sup> For example, the prevalence of primary infertility in the Middle East and north Africa, based on demographic and reproductive health surveys, has been

*Lancet Glob Health* 2019;

7: e1197–225

See [Comment](#) page e1152

\*Contributed equally

Infectious Disease Epidemiology Group, Weill Cornell Medicine-Qatar, Cornell University, Qatar Foundation-Education City, Doha, Qatar (A Smolak PhD, H Chemaitelly MSc, Prof L J Abu-Raddad PhD); Department of Communicable Diseases, HIV/Hepatitis/STIs Unit, World Health Organization Regional Office for the Eastern Mediterranean, Cairo, Egypt (J G Hermez MPH); Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland (Prof N Low PhD); Department of Healthcare Policy & Research, Weill Cornell Medicine, Cornell University, New York, NY, USA (Prof L J Abu-Raddad); and College of Health and Life Sciences, Hamad bin Khalifa University, Doha, Qatar (Prof L J Abu-Raddad)

Correspondence to: Prof Laith J Abu-Raddad, Infectious Disease Epidemiology Group, Weill Cornell Medicine-Qatar, Cornell University, Qatar Foundation-Education City, PO Box 24144, Doha, Qatar [lja2002@qatar-med.cornell.edu](mailto:lja2002@qatar-med.cornell.edu)

### Research in context

#### Evidence before this study

In a context of continuing stigma and political and sociocultural sensitivities, the Middle East and north Africa region has a dearth of epidemiological data about sexually transmitted infections. The prevalence of *Chlamydia trachomatis* and its distribution among populations at differing levels of risk of exposure remain largely unknown. A PubMed search using the search criteria ("Chlamydia"[MeSH] AND "Review" [Publication Type]) identified no systematic review and meta-analysis of regional scope for all subpopulations for this infection in the Middle East and north Africa or elsewhere.

#### Added value of this study

Using rigorous state-of-the-art methodologies with current empirical evidence, this study provided the first comprehensive epidemiological assessment of *C trachomatis* infection in the Middle East and north Africa. The study searched diverse sources of data, beyond international electronic databases, and identified a large volume of published and unpublished data, some of which now appears in the literature for the first time. The scope of evidence allowed analyses that found revealing associations relevant for the Middle East and north Africa and elsewhere. Unexpectedly, given this region's sexually

conservative norms, the study estimated a *C trachomatis* prevalence of 3% in the population at large, similar to estimates from other regions. The study also documented high *C trachomatis* prevalence levels in infertility clinic attendees and in women with miscarriage, with odds of infection three-times higher than in the general population.

#### Implications of all the available evidence

There is a substantial *C trachomatis* infection and disease burden in the Middle East and north Africa that is neglected and poorly recognised despite its social and economic toll in a region comprising 10% of the world's population. *C trachomatis* infection appears to be consistently associated with infertility and poor reproductive health outcomes in this region, yet these conditions are not linked to the possibility of an underlying infectious cause. The Middle East and north Africa is far from achieving WHO's Global Health Sector Strategy on Sexually Transmitted Infections, 2016–21. The findings of this study provide a scientific foundation to develop an evidence-informed public health response against *C trachomatis* and its burdensome sequelae. The challenge will be to implement effective targeted, culturally appropriate, and gender-specific programmes to tackle *C trachomatis* infection and improve sexual health in general.

estimated to be the highest worldwide (although that of secondary infertility seems to be the lowest).<sup>12</sup> Still, the contribution of *C trachomatis*, or other STIs, to this disease burden remains unknown. Against this background, our study aimed to characterise *C trachomatis* epidemiology in the Middle East and north Africa.

## Methods

### Search strategy and selection criteria

We did a systematic review as well as a meta-analysis and meta-regression. We followed systematic review methods proposed by the Cochrane Collaboration,<sup>13</sup> and report findings following the Preferred Reporting Items for Systematic Reviews and Meta-analyses guidelines<sup>14</sup> (appendix pp 5–6). We did exhaustive searches using PubMed and Embase, regional and national databases (WHO Index Medicus for the Eastern Mediterranean region, Iraqi Academic Scientific Journals database, and Iranian Scientific Information Database), abstract archives of International AIDS Society Conferences,<sup>15</sup> as well as country-level and international organisations' reports available through the Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project database.<sup>10,11</sup>

Our searches were done up to March 13, 2019, using broad search terms (MeSH/Emtree terms exploded to cover all subheadings and free-text terms) with no language or year restrictions. The appendix (p 7) summarises the search criteria and search terms used. The Middle East and north Africa were defined as 23 countries extending from Morocco in the west to Pakistan in the east (appendix p 8).

This definition for the Middle East and north Africa follows earlier convention applied in HIV and hepatitis C research,<sup>10,16–22</sup> and is based on definitions by WHO, UNAIDS, and the World Bank.

Search results were checked for duplicates using Endnote (version 8.2). We screened titles and abstracts of unique citations. Full texts of citations deemed relevant or potentially relevant were retrieved for further screening by AS and HC. Any document or report including biological measures for *C trachomatis* prevalence or incidence, or both, based on primary data was eligible for inclusion. Case reports, case series, editorials, commentaries, reviews, and reports about military personnel stationed in the Middle East and north Africa, but not from these countries, were excluded. Reference lists of literature reviews and all relevant articles were hand-searched for additional eligible reports.

In this Article, the term report refers to a document (article, conference abstract, or country-level report) containing outcome measures of interest (ie, prevalence or incidence) for one or more populations, and the term study refers to details of a specific outcome measure in a specific population. Consequently, one report could contribute multiple studies and one study could be published in different reports. Duplicate study results were included only once using the most detailed report.

### Data analysis

Data from relevant reports were extracted by AS with input from LJA-R. Independent extraction was done by

See Online for appendix

HC, and discrepancies were settled by consensus, or by contacting authors. Data from non-English articles were extracted from the full text by native speakers.

We extracted all measures of current (genital or rectal), recent, and ever infection with *C trachomatis*. We stratified data according to the study population's risk of exposure to *C trachomatis* or clinical manifestations (panel). Populations were defined as per original study authors' specific population definition and inclusion criteria (such as for infertile populations or women with miscarriage). We classified women and men as symptomatic only if there was an indication for the presence of *C trachomatis*-related signs and symptoms. We subsequently synthesised data by type of assay used for *C trachomatis* detection and summarised these data using medians and ranges.

Studies applying the same assay to different biological specimens were included only once, based on a sequential order that prioritised, for women, *C trachomatis* detection in endocervical swabs, followed by vaginal and urine samples; and for men, detection in urethral swabs, followed by urine and semen samples. Studies applying nucleic acid amplification test (NAAT) and culture to the same biological specimen were included separately given our interest in studying their contribution to heterogeneity in *C trachomatis* prevalence, and in generating STI-estimation correction factors based on assay type.<sup>23–25</sup> Studies applying other antigen detection assays to the same biological specimen were included only once based on assay sensitivity (direct fluorescence and enzyme-linked immunoassays on genital samples were prioritised over Giemsa staining).

We excluded studies using tissue specimens from the upper genital tract, or including less than ten participants. We stratified the analyses by sex where relevant. Studies reporting only an overall measure for men and women were classified according to the predominant sex in the sample.

We did risk of bias and precision assessments. Informed by the Cochrane approach,<sup>13</sup> we classified studies as having low versus high risk of bias for each of three quality domains assessing rigour of sampling methodology (probability based vs non-probability based), type of *C trachomatis* ascertainment (biological assay vs other, such as self-report), and response rate ( $\geq 80\%$  response rate or  $\geq 80\%$  of target sample size reached [the latter for studies using respondent-driven sampling] vs  $< 80\%$ ). Studies with unavailable information about any given domain were classified as having unclear risk of bias for that domain. Studies were considered of higher precision if 200 participants or more underwent testing for *C trachomatis*, which was judged as an acceptable level of precision assuming a mean prevalence of 3% in the general population.

We produced forest plots to visualise estimates of prevalence and 95% CIs for each at-risk population, stratified by type of assay. Pooled average prevalence and 95% CIs were then estimated using meta-analysis for each stratum. A Freeman-Tukey type arcsine square-root

#### Panel: Definitions for at-risk population classification

##### General populations (populations at low risk)

Populations at low risk of exposure to *Chlamydia trachomatis* infection such as antenatal clinic attendees, blood donors, and pregnant women.

##### Populations at intermediate risk

Populations who presumably might have some sexual contacts with populations engaging in high sexual risk behaviour, and have therefore a higher risk of exposure to *C trachomatis* than the general population. These populations comprise prisoners, people who inject drugs, truck drivers, migrant workers, and HIV-infected individuals in a setting where the HIV epidemic is driven by injecting drug use.

##### Populations at high risk

Populations at high risk of exposure to *C trachomatis* as a consequence of specific high sexual risk behaviours such as female sex workers, men who have sex with men, male sex workers, men-to-women transgenders, and HIV-infected individuals in a setting where the HIV epidemic is driven by sexual transmission.

##### Infertility clinic attendees

Infertile women or men and their partners were included in a separate category given the potential biological link between *C trachomatis* infection and infertility.

##### Women with miscarriage

These women were included in a separate category given the potential biological link between *C trachomatis* infection and miscarriage.

##### Women with ectopic pregnancy

These women were included in a separate category given the potential biological link between *C trachomatis* infection and ectopic pregnancy.

##### Symptomatic women

Women with clinical manifestations related to *C trachomatis* infection, or suspected of having a *C trachomatis* infection such as those with vaginal discharge.


##### Symptomatic men

Men with clinical manifestations related to *C trachomatis* infection, or suspected of having a *C trachomatis* infection such as those with urethral discharge.

transformation was first applied to stabilise variances of prevalence measures.<sup>26,27</sup> Measures were then weighted using the inverse-variance method,<sup>27,28</sup> before being pooled using a DerSimonian-Laird random-effects model.<sup>29</sup> This model assumes a normal distribution for true effect sizes (ie, prevalence) across studies, which factors in sampling variation and true between-study heterogeneity.<sup>30</sup>

We did heterogeneity assessment using Cochran's  $Q$  statistic to confirm existence of heterogeneity across studies,  $I^2$  to quantify magnitude of between-study variation that is due to true differences in effect size rather than chance, and prediction interval to estimate the 95% CI of the distribution of true effect sizes.<sup>30,31</sup> We did subgroup meta-analyses whenever five studies or more were available, using the R software (version 3.4.2).<sup>32</sup>

We did random-effects meta-regression analyses to identify sources of between-study heterogeneity and estimated the magnitude of their association with prevalence. We included risk of bias and precision domains in the meta-regression analyses. We considered the



**Figure 1: Flow chart of the study selection process for the systematic review**

The systematic review focused on *Chlamydia trachomatis* incidence or prevalence, or both, in the Middle East and north Africa. \*Reasons for exclusion: 14 studies applied the same assay to different biological specimens, and prevalence was included only once based on a sequential order prioritising for women *C trachomatis* detection in endocervical swabs followed by vaginal and urine samples, and for men detection in urethral swabs followed by urine and semen samples; one study applied antigen detection assays (other than nucleic acid amplification assays and culture) to the same biological specimen, and prevalence was included only once based on assay sensitivity; two studies were conducted only on subsamples of an original sample, and the prevalence in the original sample was included in the systematic review; six studies and three reports used tissue specimens from the upper genital tract; and one study had a sample size of less than 10.

following predictors a priori: at-risk population (panel), assay type (NAAT, culture, other assays detecting current infection, serological assays detecting anti-*C trachomatis* immunoglobulins of class IgG, IgM, IgA, immunoglobulins not specified, and unclear), sampling methodology (non-probability-based sampling vs probability-based sampling), sample size (<200 vs ≥200 participants), response rate (≥80% vs <80% and unclear), year of publication, year of data collection, country (Egypt, Iran,

Pakistan, and remaining countries; Egypt, Iran, and Pakistan being the most populous in the Middle East and north Africa),<sup>33</sup> and sex (women vs men; men-to-women transgenders who were biologically males were considered as men).

Studies that assessed *C trachomatis* prevalence using different diagnostics or biomarkers were included independently. Missing values for year of data collection were imputed using data for year of publication adjusted

|   | Country | Study design    | Sampling*                   | Sample size | Sex | Study context              | Population characteristics                  | Specimen     | Assay type  | Prevalence† |
|---|---------|-----------------|-----------------------------|-------------|-----|----------------------------|---|--------------|-------------|-------------|
| Kadi et al (1990) <sup>35</sup>               | Algeria | Cross-sectional | Convenience                 | 69          | W   | Gynaecology clinic         | Gynaecology clinic attendees                | Serum        | MIF (IgG)   | 17.4%       |
| Kadi et al (1990) <sup>35</sup>               | Algeria | Cross-sectional | Simple random sampling      | 180         | W   | Hospital                   | Women seeking rubella tests                 | Serum        | MIF (IgG)   | 26.6%       |
| Abdel Monem et al (2005) <sup>36</sup>        | Egypt   | Case-control    | Convenience                 | 20          | W   | Antenatal clinic           | Pregnant women                              | Endocervical | Culture     | 15%         |
| Aboul Atta and Ibrahim (1995) <sup>37</sup>   | Egypt   | Case-control    | Convenience                 | 20          | M   | Hospital                   | Controls in STI study                       | Urethral     | DFA         | 5%          |
| Badary (1996) <sup>38</sup>                   | Egypt   | Case-control    | Convenience                 | 32          | W   | Gynaecology clinic         | Fertile women                               | Endocervical | DFA         | 12.5%       |
| Berry and El Shabrawy (1996) <sup>39</sup>    | Egypt   | Case-control    | Convenience                 | 30          | W   | Family planning clinic     | Family planning clinic attendees            | Serum        | EIA (IgG)   | 3.3%        |
| Diab (1993) <sup>40</sup>                     | Egypt   | Case-control    | Convenience                 | 30          | W   | Antenatal clinic           | Women with full-term delivery               | Serum        | EIA (IgG)   | 0           |
| Draz et al (2018) <sup>41</sup>               | Egypt   | Case-control    | Convenience                 | 14          | W   | Gynaecology clinic         | Healthy women                               | Endocervical | DFA         | 0           |
| El-Sayed et al (2002) <sup>42</sup>           | Egypt   | Cross-sectional | Convenience                 | 108         | W   | Family planning clinic     | Family planning clinic attendees            | Urine        | NAAT        | 2.8%        |
| El-Sayed et al (2002) <sup>42</sup>           | Egypt   | Cross-sectional | Convenience                 | 604         | W   | Antenatal clinic           | Antenatal clinic attendees                  | Urine        | NAAT        | 1.3%        |
| Mosbah and Nabel (2016) <sup>43</sup>         | Egypt   | Case-control    | Convenience                 | 90          | W   | Hospital                   | Pregnant women with pre-eclampsia           | Endocervical | NAAT        | 4.4%        |
| Mosbah and Nabel (2016) <sup>43</sup>         | Egypt   | Case-control    | Convenience                 | 90          | W   | Hospital                   | Normotensive pregnant women                 | Endocervical | NAAT        | 0           |
| Mousa (1990) <sup>44</sup>                    | Egypt   | Cross-sectional | Convenience                 | 50          | W   | Gynaecology clinic         | Gynaecology clinic attendees                | Endocervical | DFA         | 2%          |
| Nada et al (2015) <sup>45</sup>               | Egypt   | Case-control    | Convenience                 | 100         | W   | Gynaecology clinic         | Gynaecology clinic attendees                | Endocervical | NAAT        | 2%          |
| Sullam et al (2001) <sup>46</sup>             | Egypt   | Cross-sectional | Multistage cluster sampling | 1344        | W   | Community                  | Household survey of women                   | Endocervical | ELISA       | 4.2%        |
| Zaki (1989) <sup>47</sup>                     | Egypt   | Cross-sectional | Convenience                 | 100         | W   | Antenatal clinic           | Pregnant women                              | Endocervical | Culture     | 3%          |
| Ahmadi et al (2016a) <sup>48</sup>            | Iran    | Case-control    | Convenience                 | 109         | W   | Family planning clinic     | Family planning clinic attendees            | Endocervical | NAAT        | 11.9%       |
| Ahmadi et al (2018) <sup>49</sup>             | Iran    | Case-control    | Convenience                 | 165         | M   | Clinic                     | Fertile men                                 | Semen        | NAAT        | 0.6%        |
| Ahmadnia et al (2016) <sup>50</sup>           | Iran    | Cross-sectional | Stratified cluster sampling | 4274        | W   | Primary health-care centre | Primary health-care centre clinic attendees | Endocervical | Culture     | 1%          |
| Badami and Salari (2001) <sup>51</sup>        | Iran    | Cross-sectional | Convenience                 | 250         | W   | Family planning clinic     | Family planning clinic attendees            | Serum        | DFA         | 0.8%        |
| Badami and Salari (2001) <sup>51</sup>        | Iran    | Cross-sectional | Convenience                 | 250         | W   | Family planning clinic     | Family planning clinic attendees            | Serum        | Unclear     | 3.2%        |
| Baghchesaraei et al (2011) <sup>52</sup>      | Iran    | Cross-sectional | Convenience                 | 328         | W   | Gynaecology clinic         | Gynaecology clinic attendees                | Serum        | EIA (IgM)   | 10.3%       |
| Bagheri et al (2018) <sup>53</sup>            | Iran    | Case-control    | Convenience                 | 60          | W   | Fertility centre           | Pregnant women                              | Vaginal      | NAAT        | 0           |
| Bagheri et al (2018) <sup>53</sup>            | Iran    | Case-control    | Convenience                 | 60          | W   | Fertility centre           | Pregnant women                              | Serum        | ELISA (IgA) | 6.7%        |
| Bagheri et al (2018) <sup>53</sup>            | Iran    | Case-control    | Convenience                 | 60          | W   | Fertility centre           | Pregnant women                              | Serum        | ELISA (IgG) | 1.7%        |
| Behrooz (2001) <sup>54</sup>                  | Iran    | Cross-sectional | Convenience                 | 400         | W   | Antenatal clinic           | Pregnant women                              | Unclear      | DFA         | 2.8%        |
| Chamani-Tabriz et al (2008) <sup>55</sup>     | Iran    | Cross-sectional | Convenience                 | 991         | W   | Community                  | Married women                               | Urine        | NAAT        | 12.8%       |
| Cheraghi et al (2014) <sup>56</sup>           | Iran    | Cross-sectional | Convenience                 | 1448        | W   | Health centres             | Non-pregnant women                          | Endocervical | Unclear     | 0.2%        |
| Dehghan et al (2017) <sup>57</sup>            | Iran    | Case-control    | Convenience                 | 250         | W   | Antenatal clinic           | Antenatal clinic attendees                  | Urine        | NAAT        | 0           |
| Dehghan et al (2017) <sup>57</sup>            | Iran    | Case-control    | Convenience                 | 250         | W   | Antenatal clinic           | Antenatal clinic attendees                  | Serum        | EIA (IgA)   | 0           |
| Dehghan et al (2017) <sup>57</sup>            | Iran    | Case-control    | Convenience                 | 250         | W   | Antenatal clinic           | Antenatal clinic attendees                  | Serum        | EIA (IgM)   | 0           |
| Dehghan et al (2017) <sup>57</sup>            | Iran    | Case-control    | Convenience                 | 250         | W   | Antenatal clinic           | Antenatal clinic attendees                  | Serum        | EIA (IgG)   | 12.8%       |
| Goshayeshi et al (2015) <sup>58</sup>         | Iran    | Case-control    | Convenience                 | 30          | W   | Fertility centre           | Fertile women                               | Endocervical | NAAT        | 3.3%        |
| Haghighi Hasanabad et al (2013) <sup>59</sup> | Iran    | Cross-sectional | Convenience                 | 399         | W   | Antenatal clinic           | Pregnant adolescents                        | Unclear      | NAAT        | 12.3%       |
| Jahromi et al (2010) <sup>60</sup>            | Iran    | Case-control    | Convenience                 | 200         | W   | Gynaecology clinic         | Women with full-term delivery               | Endocervical | DFA         | 5.2%        |
| Javanmard et al (2018) <sup>61</sup>          | Iran    | Cross-sectional | Convenience                 | 210         | W   | Gynaecology clinic         | Women undergoing routine pap smear          | Endocervical | NAAT        | 11.4%       |

(Table 1 continues on next page)

|   | Country | Study design    | Sampling*              | Sample size | Sex | Study context          | Population characteristics                 | Specimen     | Assay type      | Prevalence† |
|---|---------|-----------------|------------------------|-------------|-----|------------------------|--|--------------|-----------------|-------------|
| (Continued from previous page)                                |         |                 |                        |             |     |                        |  |              |                 |             |
| Joolayi et al (2017) <sup>62</sup>                            | Iran    | Case-control    | Convenience            | 125         | W   | Hospital               | Pregnant women                             | Vaginal      | NAAT            | 1.6%        |
| Joolayi et al (2017) <sup>62</sup>                            | Iran    | Case-control    | Convenience            | 125         | W   | Hospital               | Pregnant women                             | Serum        | ELISA (IgM)     | 1.6%        |
| Joolayi et al (2017) <sup>62</sup>                            | Iran    | Case-control    | Convenience            | 125         | W   | Hospital               | Pregnant women                             | Serum        | ELISA (IgG)     | 0           |
| Kajbaf and Gholamnezhad (1998) <sup>63</sup>                  | Iran    | Case-control    | Convenience            | 50          | W   | Antenatal clinic       | Antenatal clinic attendees                 | Endocervical | DFA             | 4%          |
| Kajbaf and Gholamnezhad (1998) <sup>63</sup>                  | Iran    | Case-control    | Convenience            | 50          | W   | Antenatal clinic       | Antenatal clinic attendees                 | Serum        | ELISA (IgG)     | 6%          |
| Kamyabi (2009) <sup>64</sup>                                  | Iran    | Case-control    | Convenience            | 35          | W   | Gynaecology clinic     | Pregnant women                             | Serum        | ELISA (IgG)     | 20%         |
| Khezerdoust et al (2009) <sup>65</sup>                        | Iran    | Cross-sectional | Convenience            | 1114        | W   | Antenatal clinic       | Pregnant women                             | Serum        | ELISA (IgG)     | 2.9%        |
| Marashi et al (2014) <sup>66</sup>                            | Iran    | Case-control    | Convenience            | 200         | W   | Antenatal clinic       | Pregnant women                             | Endocervical | DFA             | 3.5%        |
| Marashi et al (2014) <sup>66</sup>                            | Iran    | Case-control    | Convenience            | 200         | W   | Antenatal clinic       | Pregnant women                             | Endocervical | NAAT            | 8.7%        |
| Ministry of Health and Medical Education (2008) <sup>67</sup> | Iran    | Case-control    | Convenience            | 41          | W   | Clinic                 | Fertile women                              | Serum        | ELISA (IgG)     | 2.4%        |
| Meidani (2009) <sup>68</sup>                                  | Iran    | Cross-sectional | Simple random sampling | 140         | M   | Laboratory             | Men—premarital or pre-employment screening | Urine        | NAAT            | 0.7%        |
| Ministry of Health and Medical Education (2008) <sup>67</sup> | Iran    | Cross-sectional | Convenience            | 70          | W   | Antenatal clinic       | Pregnant women                             | Serum        | ELISA (unclear) | 4.3%        |
| Ministry of Health and Medical Education (2008) <sup>67</sup> | Iran    | Case-control    | Convenience            | 250         | W   | Family planning clinic | Healthy women                              | Endocervical | DFA             | 0.8%        |
| Mousavi et al (2014) <sup>69</sup>                            | Iran    | Case-control    | Convenience            | 104         | W   | Antenatal clinic       | Antenatal clinic attendees                 | Endocervical | NAAT            | 5.8%        |
| Pourabbas et al (2018) <sup>70</sup>                          | Iran    | Cross-sectional | Convenience            | 239         | W   | Hospital               | Pregnant women                             | Endocervical | NAAT            | 15.5%       |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience            | 223         | W   | Antenatal clinic       | Pregnant women                             | Serum        | ELISA (IgM)     | 1.8%        |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience            | 223         | W   | Antenatal clinic       | Pregnant women                             | Serum        | ELISA (IgG)     | 5.0%        |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience            | 223         | W   | Antenatal clinic       | Pregnant women                             | Urine        | NAAT            | 8.5%        |
| Rohi et al (2011) <sup>72</sup>                               | Iran    | Cross-sectional | Convenience            | 91          | W   | Hospital               | Pregnant women                             | Serum        | ELISA (IgG)     | 28.6%       |
| Rostami et al (2016) <sup>73</sup>                            | Iran    | Cross-sectional | Simple random sampling | 518         | W   | Gynaecology clinic     | Gynaecology clinic attendees               | Endocervical | NAAT            | 7.1%        |
| Safdari et al (2015) <sup>74</sup>                            | Iran    | Cross-sectional | Convenience            | 70          | W   | Antenatal clinic       | Antenatal clinic attendees                 | Endocervical | NAAT            | 10%         |
| Safdari et al (2015) <sup>74</sup>                            | Iran    | Cross-sectional | Convenience            | 70          | W   | Antenatal clinic       | Antenatal clinic attendees                 | Endocervical | Culture         | 8.6%        |
| Sattari et al (2017) <sup>75</sup>                            | Iran    | Case-control    | Convenience            | 100         | W   | Antenatal clinic       | Pregnant women                             | Serum        | ELISA (IgM)     | 2%          |
| Sattari et al (2017) <sup>75</sup>                            | Iran    | Case-control    | Convenience            | 100         | W   | Antenatal clinic       | Pregnant women                             | Serum        | ELISA (IgG)     | 18%         |
| Sisakht et al (2017) <sup>76</sup>                            | Iran    | Case-control    | Convenience            | 30          | W   | Gynaecology clinic     | Women with full-term delivery              | Urine        | NAAT            | 4.7%        |
| Yeganeh et al (2013) <sup>77</sup>                            | Iran    | Case-control    | Convenience            | 100         | M   | Urology clinic         | Asymptomatic men                           | Urine        | NAAT            | 4%          |
| Zahirmia et al (2018) <sup>78</sup>                           | Iran    | Cross-sectional | Convenience            | 76          | W   | Gynaecology clinic     | Pregnant women                             | Vaginal      | NAAT            | 10.5%       |
| Abdulkhudher et al (2014) <sup>79</sup>                       | Iraq    | Case-control    | Convenience            | 40          | W   | Antenatal clinic       | Women with full-term delivery              | Serum        | ELISA (IgM)     | 0           |
| Abdulkhudher et al (2014) <sup>79</sup>                       | Iraq    | Case-control    | Convenience            | 40          | W   | Antenatal clinic       | Women with full-term delivery              | Serum        | ELISA (IgG)     | 7.5%        |
| Abdul-Karim et al (2009) <sup>80</sup>                        | Iraq    | Cross-sectional | Convenience            | 198         | W   | Hospital               | Women with full-term delivery              | Serum        | ELISA (IgG)     | 13.7%       |
| Abdullah (2012) <sup>81</sup>                                 | Iraq    | Case-control    | Convenience            | 24          | W   | Hospital               | Pregnant women                             | Serum        | ELISA (IgM)     | 0           |
| Abdullah (2012) <sup>81</sup>                                 | Iraq    | Case-control    | Convenience            | 24          | W   | Hospital               | Pregnant women                             | Serum        | ELISA (IgG)     | 8.3%        |
| Ahmed (2008) <sup>82</sup>                                    | Iraq    | Case-control    | Convenience            | 30          | W   | Hospital               | Women with full-term delivery              | Serum        | ELISA (unclear) | 0           |
| Al-Hamdani et al (2010) <sup>83</sup>                         | Iraq    | Case-control    | Convenience            | 17          | W   | Hospital               | Pregnant women                             | Serum        | ELISA (IgM)     | 14.0%       |

(Table 1 continues on next page)



|  | Country | Study design    | Sampling*                  | Sample size | Sex | Study context              | Population characteristics               | Specimen     | Assay type     | Prevalence† |
|--|---------|-----------------|----------------------------|-------------|-----|----------------------------|--|--------------|----------------|-------------|
| (Continued from previous page)                 |         |                 |                            |             |     |                            |  |              |                |             |
| Al-Hamdani et al (2010) <sup>83</sup>          | Iraq    | Case-control    | Convenience                | 17          | W   | Hospital                   | Pregnant women                           | Serum        | ELISA (IgG)    | 14.0%       |
| Al-Hamdani et al (2010) <sup>83</sup>          | Iraq    | Case-control    | Convenience                | 17          | W   | Hospital                   | Pregnant women                           | Serum        | ELISA (IgA)    | 40.4%       |
| Al-Husseinei et al (2009) <sup>84</sup>        | Iraq    | Case-control    | Convenience                | 100         | W   | Family planning clinic     | Family planning clinic attendees         | Serum        | IFAT (unclear) | 5%          |
| Al-Husseinei et al (2009) <sup>84</sup>        | Iraq    | Case-control    | Convenience                | 100         | W   | Family planning clinic     | Family planning clinic attendees         | Endocervical | ELFA           | 4%          |
| Ali and Al-Kazaz (2018) <sup>85</sup>          | Iraq    | Case-control    | Convenience                | 13          | M   | Clinic                     | Fertile men                              | Semen        | NAAT           | 0           |
| Alkhafaf (2013) <sup>86</sup>                  | Iraq    | Case-control    | Convenience                | 122         | W   | Hospital                   | Married women                            | Serum        | ELISA (IgG)    | 4.1%        |
| Alkhafaf (2013) <sup>86</sup>                  | Iraq    | Case-control    | Convenience                | 168         | W   | Hospital                   | Unmarried woman                          | Serum        | ELISA (IgG)    | 2.4%        |
| Hwaid et al (2013) <sup>87</sup>               | Iraq    | Case-control    | Simple random sampling     | 91          | W   | Antenatal clinic           | Pregnant women                           | Serum        | ELISA (IgG)    | 5.5%        |
| Hwaid et al (2013) <sup>87</sup>               | Iraq    | Case-control    | Simple random sampling     | 91          | W   | Antenatal clinic           | Pregnant women                           | Serum        | ELISA (IgM)    | 3.3%        |
| Ismail and Ali (2012) <sup>88</sup>            | Iraq    | Case-control    | Convenience                | 50          | W   | Laboratories               | General population women                 | Serum        | ELISA (IgM)    | 10%         |
| Ismail and Ali (2012) <sup>88</sup>            | Iraq    | Case-control    | Convenience                | 50          | W   | Laboratories               | General population women                 | Serum        | ELISA (IgG)    | 4%          |
| Ismail and Ali (2012) <sup>88</sup>            | Iraq    | Case-control    | Convenience                | 50          | W   | Laboratories               | General population women                 | Serum        | ELISA (IgA)    | 2%          |
| Mohammed et al (2012) <sup>89</sup>            | Iraq    | Case-control    | Convenience                | 23          | W   | Gynaecology clinic         | Pregnant women                           | Serum        | ELISA (IgM)    | 0           |
| Mohammed et al (2017) <sup>90</sup>            | Iraq    | Case-control    | Convenience                | 20          | W   | Gynaecology clinic         | Gynaecology clinic attendees             | Endocervical | NAAT           | 0           |
| Mohammed et al (2017) <sup>90</sup>            | Iraq    | Case-control    | Convenience                | 20          | W   | Gynaecology clinic         | Gynaecology clinic attendees             | Serum        | ELISA (IgG)    | 0           |
| Yahya and Al-Siraj (2009) <sup>91</sup>        | Iraq    | Cross-sectional | Convenience                | 296         | M   | Laboratory                 | Fertile men                              | Serum        | Culture        | 0           |
| Abusarah et al (2013) <sup>92</sup>            | Jordan  | Case-control    | Convenience                | 61          | M   | Urology clinics            | Fertile men                              | Urine        | NAAT           | 1.6%        |
| Al-Ramahi et al (2008) <sup>93</sup>           | Jordan  | Case-control    | Convenience                | 146         | W   | Gynaecology clinic         | Gynaecology clinic attendees             | Endocervical | NAAT           | 0.7%        |
| As'ad (2004) <sup>94</sup>                     | Jordan  | Cross-sectional | Convenience                | 144         | W   | Family planning clinic     | Asymptomatic women                       | Vaginal      | NAAT           | 0           |
| Awwad et al (2003) <sup>95</sup>               | Jordan  | Case-control    | Convenience                | 61          | M   | Urology clinic             | Non-urethritis patients                  | Urine        | NAAT           | 0           |
| Awwad et al (2003) <sup>95</sup>               | Jordan  | Case-control    | Convenience                | 39          | W   | Urology clinic             | Non-urethritis patients                  | Urine        | NAAT           | 0           |
| Mahafzah et al (2008) <sup>96</sup>            | Jordan  | Cross-sectional | Convenience                | 186         | W   | Gynaecology clinic         | Family planning clinic attendees         | Endocervical | NAAT           | 0.5%        |
| Jordan Ministry of Health (2004) <sup>97</sup> | Jordan  | Cross-sectional | Convenience                | 213         | W   | Hospital                   | Asymptomatic women                       | Endocervical | NAAT           | 0.5%        |
| Al-Awadhi et al (2018) <sup>98</sup>           | Kuwait  | Cross-sectional | Convenience                | 65 338      | W   | Laboratory                 | Women undergoing pap smear 1997–2005     | Endocervical | Unclear        | 0.1%        |
| Al-Awadhi et al (2018) <sup>98</sup>           | Kuwait  | Cross-sectional | Convenience                | 56 105      | W   | Laboratory                 | Women undergoing pap smear 2006–14       | Endocervical | Unclear        | 0.04%       |
| Al-Sweih et al (2011) <sup>99</sup>            | Kuwait  | Cross-sectional | Convenience                | 5938        | W   | Primary health-care centre | Kuwaiti women                            | Vaginal      | NAAT           | 1.9%        |
| Al-Sweih et al (2011) <sup>99</sup>            | Kuwait  | Cross-sectional | Convenience                | 2601        | W   | Primary health-care centre | Expatriate women                         | Vaginal      | NAAT           | 2.3%        |
| Al-Sweih et al (2012) <sup>100</sup>           | Kuwait  | Case-control    | Convenience                | 188         | M   | Gynaecology clinic         | Fertile men                              | Semen        | NAAT           | 3.7%        |
| Deeb et al (2003) <sup>101</sup>               | Lebanon | Cross-sectional | Multistage random sampling | 506         | W   | Community                  | Ever-married women                       | Endocervical | ELISA          | 0           |
| Hancali et al (2015) <sup>102</sup>            | Morocco | Cross-sectional | Convenience                | 760         | W   | Family planning clinic     | Family planning clinic attendees in 1999 | Unclear      | NAAT           | 4.0%        |
| Hancali et al (2015) <sup>102</sup>            | Morocco | Cross-sectional | Convenience                | 256         | W   | Family planning clinic     | Family planning clinic attendees in 2011 | Unclear      | NAAT           | 4.4%        |

(Table 1 continues on next page)

(Table 1 continues on next page)

|  | Country      | Study design                 | Sampling*                             | Sample size | Sex | Study context                               | Population characteristics                            | Specimen     | Assay type     | Prevalence† |
|--|--------------|------------------------------|---------------------------------------|-------------|-----|---|---|--------------|----------------|-------------|
| (Continued from previous page)   |              |                              |                                       |             |     |   |   |              |                |             |
| Hulstein et al (2018) <sup>103</sup>   | Morocco      | Cross-sectional              | Simple random sampling                | 163         | M   | Community                                   | General population men                                | Serum        | IFAT (IgG)     | 31.3%       |
| Hulstein et al (2018) <sup>103</sup>   | Morocco      | Cross-sectional              | Simple random sampling                | 174         | W   | Community                                   | General population women                              | Serum        | IFAT (IgG)     | 37.9%       |
| Morocco Ministry of Health (2001) <sup>104</sup>   | Morocco      | Cross-sectional              | Convenience                           | 323         | W   | Antenatal clinic                            | Pregnant women  | Urine        | NAAT           | 2.7%        |
| Morocco Ministry of Health (2001) <sup>104</sup>   | Morocco      | Cross-sectional              | Convenience                           | 518         | W   | Family planning clinic                      | Family planning clinic attendees                      | Urine        | NAAT           | 5.2%        |
| The Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project (2017) <sup>105</sup> | Morocco      | Cross-sectional              | Convenience                           | 252         | W   | Antenatal clinic                            | Pregnant women  | Unclear      | NAAT           | 3.6%        |
| The Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project (2017) <sup>105</sup> | Morocco      | Cross-sectional              | Convenience                           | 537         | W   | Family planning clinic                      | Family planning clinic attendees                      | Unclear      | NAAT           | 3%          |
| Radouani et al (1998) <sup>106</sup>   | Morocco      | Case-control                 | Convenience                           | 81          | W   | Hospital                                    | Pregnant women  | Serum        | MIF (unclear)  | 14.8%       |
| Radouani et al (1998) <sup>106</sup>   | Morocco      | Case-control                 | Convenience                           | 200         | M   | Hospital                                    | Blood donors  | Serum        | MIF (unclear)  | 4.5%        |
| Takourt et al (1995) <sup>107</sup>  | Morocco      | Case-control                 | Convenience                           | 200         | M   | Hospital                                    | Blood donors  | Serum        | MIF (unclear)  | 5.0%        |
| Takourt et al (1995) <sup>107</sup>  | Morocco      | Case-control                 | Convenience                           | 200         | W   | Hospital                                    | Blood donors  | Serum        | MIF (unclear)  | 10.0%       |
| Mir et al (2009) <sup>108</sup>  | Pakistan     | Cross-sectional              | Multistage systematic random sampling | 2383        | M   | Community                                   | General population men                                | Urine        | NAAT           | 0           |
| Wasti et al (1997) <sup>109</sup>  | Pakistan     | Cross-sectional              | Convenience                           | 300         | W   | Antenatal clinic and family planning clinic | Antenatal clinic and family planning clinic attendees | Endocervical | DFA            | 5.3%        |
| Al-Thani et al (2013) <sup>110</sup>   | Qatar        | Cross-sectional              | Convenience                           | 133         | W   | Primary health-care centre                  | Qatari women  | Endocervical | NAAT           | 5.3%        |
| Al-Thani et al (2013) <sup>110</sup>   | Qatar        | Cross-sectional              | Convenience                           | 218         | W   | Primary health-care centre                  | Non-Qatari women                                      | Endocervical | NAAT           | 5.5%        |
| Alzahrani et al (2010) <sup>111</sup>  | Saudi Arabia | Cross-sectional              | Simple random sampling                | 95          | W   | Antenatal clinic                            | Pregnant women  | Endocervical | ELISA          | 10.5%       |
| Awad et al (2013) <sup>112</sup>   | Saudi Arabia | Cross-sectional              | Convenience                           | 144         | W   | Gynaecology clinic                          | Antenatal clinic attendees                            | Urine        | NAAT           | 11.1%       |
| Bashi (1987) <sup>113</sup>  | Saudi Arabia | Cross-sectional              | Convenience                           | 100         | W   | Primary health-care centre                  | Primary health-care centre attendees                  | Serum        | MIF (IgG)      | 0           |
| Bashi (1987) <sup>113</sup>  | Saudi Arabia | Cross-sectional              | Convenience                           | 100         | M   | Primary health-care centre                  | Primary health-care centre attendees                  | Serum        | MIF (IgG)      | 2%          |
| Ghazi et al (2006) <sup>114</sup>  | Saudi Arabia | Cross-sectional              | Simple random sampling                | 1600        | W   | Antenatal clinic                            | Saudi pregnant women                                  | Serum        | ELISA (IgG)    | 8.7%        |
| Ghazi et al (2006) <sup>114</sup>  | Saudi Arabia | Cross-sectional              | Simple random sampling                | 1460        | W   | Antenatal clinic                            | Saudi pregnant women                                  | Serum        | ELISA (IgM)    | 1.5%        |
| Hossain (1988) <sup>115</sup>  | Saudi Arabia | Cross-sectional              | Convenience                           | 112         | M   | Hospital                                    | Blood donors  | Serum        | MIF (IgM)      | 0           |
| Hossain (1988) <sup>115</sup>  | Saudi Arabia | Cross-sectional              | Convenience                           | 112         | M   | Hospital                                    | Blood donors  | Serum        | MIF (IgG)      | 1.8%        |
| Kamel (2013) <sup>116</sup>  | Saudi Arabia | Randomised controlled trial‡ | Convenience                           | 100         | W   | Antenatal clinic                            | Antenatal clinic attendees                            | Serum        | ELISA (IgG)    | 4.0%        |
| Massoud et al (1991) <sup>117</sup>  | Saudi Arabia | Case-control                 | Convenience                           | 100         | W   | Hospital                                    | Asymptomatic women                                    | Serum        | IFAT (unclear) | 0           |
| Massoud et al (1991) <sup>117</sup>  | Saudi Arabia | Case-control                 | Convenience                           | 100         | M   | Hospital                                    | Asymptomatic men                                      | Serum        | IFAT (unclear) | 2.0%        |
| Ismail et al (1990) <sup>118</sup>   | Somalia      | Cross-sectional              | Convenience                           | 194         | W   | Community                                   | Women   | Endocervical | EIA            | 12.4%       |

(Table 1 continues on next page)



|  | Country              | Study design    | Sampling*                   | Sample size | Sex | Study context    | Population characteristics                      | Specimen                  | Assay type  | Prevalence† |
|--|----------------------|-----------------|-----------------------------|-------------|-----|------------------|---|---------------------------|-------------|-------------|
| (Continued from previous page)           |                      |                 |                             |             |     |                  |   |                           |             |             |
| Ismail et al (1990) <sup>118</sup>       | Somalia              | Cross-sectional | Convenience                 | 189         | M   | Community        | Men   | Urethral                  | EIA         | 6%          |
| Nur et al (2000) <sup>119</sup>          | Somalia              | Cross-sectional | Convenience                 | 54          | M   | Hospital         | Blood donors                                    | Serum                     | EIA (IgG)   | 22.2%       |
| WHO (2005a) <sup>120</sup>               | Somalia              | Cross-sectional | Convenience                 | 4723        | W   | Antenatal clinic | Pregnant women                                  | Urine                     | NAAT        | 1.7%        |
| WHO (2005b) <sup>121</sup>               | Somalia              | Cross-sectional | Convenience                 | 509         | W   | Antenatal clinic | Pregnant women                                  | Urine                     | NAAT        | 1.4%        |
| Ahmed et al (2018) <sup>122</sup>        | Sudan                | Case-control    | Convenience                 | 93          | W   | Hospital         | Healthy pregnant women                          | Serum                     | ELISA (IgG) | 0           |
| Ahmed et al (2018) <sup>122</sup>        | Sudan                | Case-control    | Convenience                 | 93          | W   | Hospital         | Pregnant women with pre-eclampsia               | Serum                     | ELISA (IgG) | 0           |
| Almroth et al (2005) <sup>123</sup>      | Sudan                | Case-control    | Convenience                 | 139         | W   | Antenatal clinic | Antenatal clinic attendees                      | Serum                     | EIA (IgG)   | 3.6%        |
| Ortashi et al (2004) <sup>124</sup>      | Sudan                | Cross-sectional | Convenience                 | 151         | W   | Antenatal clinic | Pregnant women                                  | Endocervical and urethral | EIA         | 19.9%       |
| Alkayer et al (2017) <sup>125</sup>      | Syria                | Case-control    | Convenience                 | 21          | W   | Hospital         | Pregnant women                                  | Serum                     | ELISA (IgG) | 4.7%        |
| Ghazal-Aswad et al (2006) <sup>126</sup> | United Arab Emirates | Cross-sectional | Multistage cluster sampling | 727         | W   | Clinics          | Primary health-care centre and clinic attendees | Endocervical              | EIA         | 2.5%        |

DFA=direct fluorescent assay. EIA=enzyme immunoassay. ELFA=enzyme-linked fluorescence assay. IFAT=indirect fluorescent antibody test. M=men or sample predominantly of men. MIF=micro-immunofluorescence. NAAT=nucleic acid amplification test. STI=sexually transmitted infection. W=women or sample predominantly of women. \*Non-probability sampling refers to a sampling method in which the data collection process does not allow individuals to have equal chance of being selected; an example is convenience sampling for which individuals are selected on the basis of ease of accessibility (first-come first-served basis).<sup>127,128</sup> Probability-based sampling refers to a sampling method in which data collection process is based on a random selection of study participants; an example is random sampling from a sampling frame.<sup>128</sup> Another example of probability-based sampling is respondent-driven sampling, which is a sampling method specifically designed to sample hard-to-reach populations and is based on chain referral with the probability of selection calculated at each step in the network to produce adjusted prevalence estimates.<sup>129</sup> †The decimal places of the prevalence measures are as reported in the original report, but prevalence figures with more than one decimal place were rounded to one decimal place. ‡The extracted prevalence measure is for the baseline measurement.

**Table 1: Studies reporting *Chlamydia trachomatis* prevalence in general populations in the Middle East and north Africa**

**Table 1: Studies reporting *Chlamydia trachomatis* prevalence in general populations in the Middle East and north Africa**

by the median difference between year of publication and year of data collection (for studies with complete information). We did meta-regression diagnostics.

Factors associated with prevalence at  $p \leq 0.20$  in univariable analysis were eligible for inclusion in the multivariable analysis. In the multivariable model, a  $p \leq 0.05$  for any given factor indicated strong evidence for an association with *C trachomatis* prevalence. We did meta-regression models using Stata/SE (version 14).<sup>34</sup>

### Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the article. The corresponding author had full access to all the data in the study and had the final responsibility for the decision to submit for publication.

### Results

The search identified a total of 1531 citations: 509 through PubMed, 557 through Embase, and 465 through regional and national databases. Of these citations, 302 reports underwent full-text screening after excluding duplicates and screening titles and abstracts. During full-text screening, 228 eligible reports were identified, and 74 were excluded for reasons outlined in figure 1. Hand-searching of reference lists of relevant reports and reviews yielded 14 additional eligible reports. 16 country-level reports were further identified through the Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project database. Three reports were subsequently excluded. In total, 255 reports contributing 552 prevalence measures

or studies met the eligibility criteria for inclusion, but no incidence measures were identified.

Evidence covered 20 (87%) of 23 countries, encompassing a total of 256 769 *C trachomatis* test results (tables 1 and 2; appendix pp 9–14). Iran contributed the largest number of measures or studies ( $n=176$ ), followed by Egypt ( $n=89$ ), Iraq ( $n=72$ ), Saudi Arabia ( $n=45$ ), Pakistan ( $n=42$ ), and Morocco ( $n=32$ ). Most studies assessed current infection ( $n=318$ ), whereas the rest reported different serological measures ( $n=211$ ), such as ever infection (anti-*C trachomatis* IgG;  $n=117$ ). Details of *C trachomatis* testing protocol were specified in 424 (77%) of 552 studies; 320 (75%) of the 424 used commercial assays, 62 (15%) used in-house validated tests, 29 (7%) used culture, and 13 (3%) used a non-validated in-house test.

In general populations ( $n=137$ ), prevalence of current genital infection ranged from 0 to 19.9% with a median of 3.0%, whereas ever infection prevalence ranged from 0 to 37.9% with a median of 4.7% (tables 1 and 3).

In populations at high risk ( $n=40$ ), current infection prevalence in female sex workers ( $n=20$ ) ranged from 0.9% to 72.9% with a median of 8.4%, whereas ever infection prevalence ranged from 19.8% to 100% with a median of 90.0% (tables 2 and 3). In men who have sex with men (including male sex workers and male-to-female transgenders;  $n=20$ ), current infection prevalence ranged from 0 to 8.8% with a median of 1.2% for genital infections and from 3.6% to 18.3% with a median of 6.3% for rectal infections, but no ever infection measure was identified.

|  | Country  | Study design    | Sampling*                  | Sample size | Sex | Study context  | Population characteristics                         | Specimen                 | Assay type     | Prevalence† |
|--|----------|-----------------|----------------------------|-------------|-----|----------------|--|--------------------------|----------------|-------------|
| <b>Populations at high risk</b>                  |          |                 |                            |             |     |                |  |                          |                |             |
| Kadi et al (1990) <sup>35</sup>                  | Algeria  | Cross-sectional | Convenience                | 44          | W   | Community      | Female sex workers                                 | Serum                    | MIF (IgG)      | 100%        |
| El-Sayed et al (2002) <sup>42</sup>              | Egypt    | Cross-sectional | Convenience                | 52          | W   | Community      | Female sex workers                                 | Urine                    | NAAT           | 7.7%        |
| El-Sayed et al (2002) <sup>42</sup>              | Egypt    | Cross-sectional | Convenience                | 80          | M   | Community      | Men who have sex with men                          | Urine                    | NAAT           | 8.8%        |
| Darougar et al (1983) <sup>130</sup>             | Iran     | Cross-sectional | Convenience                | 116         | W   | Community      | Female sex workers                                 | Endocervical             | Culture        | 6.9%        |
| Darougar et al (1983) <sup>130</sup>             | Iran     | Cross-sectional | Convenience                | 154         | W   | Community      | Female sex workers                                 | Serum                    | MIF (IgM)      | 29.2%       |
| Darougar et al (1983) <sup>130</sup>             | Iran     | Cross-sectional | Convenience                | 154         | W   | Community      | Female sex workers                                 | Serum                    | MIF (IgG)      | 94.2%       |
| Kassaian et al (2012) <sup>131</sup>             | Iran     | Cross-sectional | Convenience                | 91          | W   | Mixed          | Female sex workers                                 | Serum                    | ELISA (IgG)    | 19.8%       |
| Kazerooni et al (2014) <sup>132</sup>            | Iran     | Cross-sectional | Respondent-driven sampling | 278         | W   | Community      | Female sex workers                                 | Vaginal                  | NAAT           | 9%          |
| Mirzazadeh et al (2016) <sup>133</sup>           | Iran     | Cross-sectional | Convenience                | 1337        | W   | Community      | Female sex workers                                 | Vaginal                  | NAAT           | 6%          |
| Al-Hussein et al (2009) <sup>84</sup>            | Iraq     | Case-control    | Convenience                | 30          | W   | STI clinic     | Women with multiple partners                       | Endocervical             | ELFA           | 30%         |
| Al-Hussein et al (2009) <sup>84</sup>            | Iraq     | Case-control    | Convenience                | 30          | W   | STI clinic     | Women with multiple partners                       | Serum                    | IFAT (unclear) | 36.7%       |
| Bellaji et al (2017) <sup>134</sup>              | Morocco  | Cross-sectional | Convenience                | 519         | W   | NGOs           | Female sex workers                                 | Endocervical and vaginal | NAAT           | 20.7%       |
| Morocco Ministry of Health (2008) <sup>135</sup> | Morocco  | Cross-sectional | Convenience                | 141         | W   | STI clinic     | Female sex workers                                 | Endocervical and urine   | NAAT           | 22.7%       |
| Morocco Ministry of Health (2011) <sup>136</sup> | Morocco  | Cross-sectional | Respondent-driven sampling | 368         | W   | Community      | Female sex workers in Agadir                       | Endocervical             | NAAT           | 22.4%       |
| Morocco Ministry of Health (2015) <sup>137</sup> | Morocco  | Cross-sectional | Respondent-driven sampling | 247         | M   | Community      | Men who have sex with men in Agadir                | Urine                    | NAAT           | 5.4%        |
| Morocco Ministry of Health (2015) <sup>137</sup> | Morocco  | Cross-sectional | Respondent-driven sampling | 252         | M   | Community      | Men who have sex with men in Marrakech             | Urine                    | NAAT           | 6.5%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 426         | W   | Community      | Female sex workers in Rawalpindi                   | Endocervical             | NAAT           | 1.7%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 107         | W   | Community      | Female sex workers in Abbottabad                   | Endocervical             | NAAT           | 0.9%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 195         | M   | Community      | Male sex workers in Rawalpindi (Bantha)            | Urine                    | NAAT           | 0           |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 195         | M   | Community      | Male sex workers in Rawalpindi (Bantha)            | Rectal                   | NAAT           | 4.7%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 364         | M   | Community      | Male sex workers in Rawalpindi (Khotki)            | Urine                    | NAAT           | 0           |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 364         | M   | Community      | Male sex workers in Rawalpindi (Khotki)            | Rectal                   | NAAT           | 3.6%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 253         | M   | Community      | Male sex workers in Rawalpindi (Khusra)            | Urine                    | NAAT           | 0           |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 253         | M   | Community      | Male sex workers in Rawalpindi (Khusra)            | Rectal                   | NAAT           | 9.9%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 83          | M   | Community      | Male sex workers in Abbottabad (Bantha)            | Urine                    | NAAT           | 1.2%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 83          | M   | Community      | Male sex workers in Abbottabad (Bantha)            | Rectal                   | NAAT           | 4.9%        |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 20          | M   | Community      | Male sex workers in Abbottabad (Khotki and Khusra) | Urine                    | NAAT           | 0           |
| Hawkes et al (2009) <sup>138</sup>               | Pakistan | Cross-sectional | Respondent-driven sampling | 20          | M   | Community      | Male sex workers in Abbottabad (Khotki and Khusra) | Rectal                   | NAAT           | 6.3%        |
| Khan et al (2011) <sup>139</sup>                 | Pakistan | Cross-sectional | Respondent-driven sampling | 730         | W   | Community      | Female sex workers in Lahore                       | Endocervical             | NAAT           | 7.7%        |
| Osama (2017) <sup>140</sup>                      | Pakistan | Cross-sectional | Convenience                | 2531        | M   | Drop in centre | Men who have sex with men in Lahore                | Unclear                  | Unclear        | 35.2%       |
| Rehan et al (2009) <sup>141</sup>                | Pakistan | Cross-sectional | Systematic random sampling | 383         | W   | Red-light      | Female sex workers in Lahore                       | Endocervical             | NAAT           | 11%         |
| Rehan et al (2009) <sup>141</sup>                | Pakistan | Cross-sectional | Snowball                   | 348         | W   | Community      | Female sex workers in Karachi                      | Endocervical             | NAAT           | 5.2%        |

(Table 2 continues on next page)

|  | Country  | Study design    | Sampling*                          | Sample size | Sex | Study context      | Population characteristics                                 | Specimen     | Assay type  | Prevalence† |
|--|----------|-----------------|------------------------------------|-------------|-----|--------------------|--|--------------|-------------|-------------|
| (Continued from previous page)             |          |                 |                                    |             |     |                    |  |              |             |             |
| Rehan et al (2009) <sup>141</sup>          | Pakistan | Cross-sectional | Respondent-driven sampling         | 395         | M   | Community          | Male sex workers in Lahore                                 | Urethral     | NAAT        | 1.5%        |
| Rehan et al (2009) <sup>141</sup>          | Pakistan | Cross-sectional | Snowball                           | 396         | M   | Community          | Male sex workers in Karachi                                | Urethral     | NAAT        | 1.2%        |
| Rehan et al (2009) <sup>141</sup>          | Pakistan | Cross-sectional | Snowball                           | 394         | M   | Community          | Male sex workers in Karachi                                | Rectal       | NAAT        | 10.4%       |
| Rehan et al (2009) <sup>141</sup>          | Pakistan | Cross-sectional | Systematic random cluster sampling | 198         | M   | Community          | Hijras in Lahore   | Urethral     | NAAT        | 1.5%        |
| Rehan et al (2009) <sup>141</sup>          | Pakistan | Cross-sectional | Systematic random cluster sampling | 197         | M   | Community          | Hijras in Karachi  | Urethral     | NAAT        | 0           |
| Rehan et al (2009) <sup>141</sup>          | Pakistan | Cross-sectional | Systematic random cluster sampling | 197         | M   | Community          | Hijras in Karachi  | Rectal       | NAAT        | 18.3%       |
| Znazen et al (2010) <sup>142</sup>         | Tunisia  | Cross-sectional | Convenience                        | 188         | W   | Community          | Female sex workers   | Endocervical | NAAT        | 72.9%       |
| Znazen et al (2010) <sup>142</sup>         | Tunisia  | Cross-sectional | Convenience                        | 183         | W   | Community          | Female sex workers   | Serum        | MIF (IgG)   | 85.8%       |
| <b>Infertility clinic attendees</b>        |          |                 |                                    |             |     |                    |  |              |             |             |
| Abdel Aleem et al (1996) <sup>143</sup>    | Egypt    | Case-control    | Convenience                        | 144         | W   | Infertility clinic | Women with mixed infertility diagnosis                     | Serum        | ELISA (IgG) | 52%         |
| Abdel Aleem et al (1996) <sup>143</sup>    | Egypt    | Case-control    | Convenience                        | 104         | M   | Infertility clinic | Men with unclear infertility diagnosis                     | Serum        | ELISA (IgG) | 24%         |
| Abdel Monem et al (2005) <sup>36</sup>     | Egypt    | Case-control    | Convenience                        | 150         | W   | Infertility clinic | Women with unclear infertility diagnosis                   | Endocervical | Culture     | 24%         |
| Abdel Monem et al (2005) <sup>36</sup>     | Egypt    | Case-control    | Convenience                        | 150         | W   | Infertility clinic | Women with unclear infertility diagnosis                   | Endocervical | EIA         | 22.7%       |
| Abdella et al (2015) <sup>144</sup>        | Egypt    | Case-control    | Convenience                        | 50          | W   | Infertility clinic | Women with idiopathic infertility                          | Serum        | ELISA (IgM) | 4%          |
| Abdella et al (2015) <sup>144</sup>        | Egypt    | Case-control    | Convenience                        | 50          | W   | Infertility clinic | Women with idiopathic infertility                          | Serum        | ELISA (IgG) | 36%         |
| Abdella et al (2015) <sup>144</sup>        | Egypt    | Case-control    | Convenience                        | 50          | W   | Infertility clinic | Women with idiopathic infertility                          | Endocervical | NAAT        | 6%          |
| Azab and Hassouna (2008) <sup>145</sup>    | Egypt    | Cross-sectional | Convenience                        | 70          | W   | Infertility clinic | Nearly half of women with TFI                              | Serum        | ELISA (IgG) | 28.6%       |
| Badary (1996) <sup>38</sup>                | Egypt    | Case-control    | Convenience                        | 60          | W   | Infertility clinic | Women with idiopathic infertility                          | Endocervical | DFA         | 33%         |
| Berry and El Shabrawy (1996) <sup>39</sup> | Egypt    | Case-control    | Convenience                        | 70          | W   | Infertility clinic | Women with unclear infertility diagnosis                   | Serum        | EIA (IgG)   | 18.6%       |
| Elkayal et al (2015) <sup>146</sup>        | Egypt    | Case-control    | Convenience                        | 100         | W   | Infertility clinic | Women with mixed infertility diagnosis                     | Endocervical | ELISA       | 3%          |
| Elkayal et al (2015) <sup>146</sup>        | Egypt    | Case-control    | Convenience                        | 100         | W   | Infertility clinic | Women with mixed infertility diagnosis                     | Endocervical | NAAT        | 3%          |
| El Sayed et al (1997) <sup>147</sup>       | Egypt    | Cross-sectional | Convenience                        | 22          | W   | Infertility clinic | Women with TFI   | Serum        | MIF (IgG)   | 81.8%       |
| El Sayed et al (1997) <sup>147</sup>       | Egypt    | Cross-sectional | Convenience                        | 78          | W   | Infertility clinic | Women without TFI  | Serum        | MIF (IgG)   | 7.7%        |
| Inhorn and Buss (1993) <sup>148</sup>      | Egypt    | Case-control    | Convenience                        | 83          | W   | Hospital           | Majority of women without TFI                              | Unclear      | Unclear     | 33%         |
| Makled et al (2013) <sup>149</sup>         | Egypt    | Cross-sectional | Simple random sampling             | 27          | W   | Infertility clinic | Women with TFI   | Serum        | ELISA (IgG) | 85.2%       |
| Makled et al (2013) <sup>149</sup>         | Egypt    | Cross-sectional | Simple random sampling             | 51          | W   | Infertility clinic | Women without TFI  | Serum        | ELISA (IgG) | 13.7%       |
| Nada et al (2015) <sup>45</sup>            | Egypt    | Case-control    | Convenience                        | 100         | W   | Infertility clinic | Women with idiopathic infertility                          | Endocervical | NAAT        | 15%         |
| Sadek et al (1993) <sup>150</sup>          | Egypt    | Case-control    | Convenience                        | 43          | W   | Infertility clinic | Infertile women in infertile couples with sperm antibodies | Unclear      | DFA         | 18.6%       |
| Sadek et al (1993) <sup>150</sup>          | Egypt    | Case-control    | Convenience                        | 37          | W   | Infertility clinic | Women partners in infertile couples with sperm antibodies  | Unclear      | DFA         | 18.9%       |
| Sadek et al (1993) <sup>150</sup>          | Egypt    | Case-control    | Convenience                        | 62          | M   | Infertility clinic | Men partners in infertile couples with sperm antibodies    | Unclear      | DFA         | 19.4%       |

(Table 2 continues on next page)

|  | Country | Study design    | Sampling*   | Sample size | Sex | Study context      | Population characteristics                               | Specimen     | Assay type  | Prevalence† |
|--|---------|-----------------|-------------|-------------|-----|--------------------|--|--------------|-------------|-------------|
| (Continued from previous page)           |         |                 |             |             |     |                    |  |              |             |             |
| Sadek et al (1993) <sup>150</sup>        | Egypt   | Case-control    | Convenience | 18          | M   | Infertility clinic | Infertile men in infertile couples with sperm antibodies | Unclear      | DFA         | 22.2%       |
| Siam and Hefzy (2012) <sup>151</sup>     | Egypt   | Case-control    | Convenience | 90          | W   | Gynaecology clinic | Women with idiopathic infertility                        | Serum        | ELISA (IgG) | 20%         |
| Siam and Hefzy (2012) <sup>151</sup>     | Egypt   | Case-control    | Convenience | 90          | W   | Gynaecology clinic | Women with idiopathic infertility                        | Urine        | NAAT        | 4.4%        |
| Younis et al (2000) <sup>152</sup>       | Egypt   | Cross-sectional | Convenience | 30          | W   | Infertility clinic | Women with TFI   | Serum        | MIF (IgG)   | 46.7%       |
| Younis et al (2000) <sup>152</sup>       | Egypt   | Cross-sectional | Convenience | 14          | W   | Infertility clinic | Women without TFI  | Serum        | MIF (IgG)   | 50.0%       |
| Zaitun and Zaitoun (1990) <sup>153</sup> | Egypt   | Cross-sectional | Convenience | 20          | W   | Infertility clinic | Women with TFI   | Serum        | Unclear     | 25%         |
| Zaitun and Zaitoun (1990) <sup>153</sup> | Egypt   | Cross-sectional | Convenience | 30          | W   | Infertility clinic | Women without TFI  | Serum        | Unclear     | 3.3%        |
| Zaki (1989) <sup>47</sup>                | Egypt   | Cross-sectional | Convenience | 100         | W   | Infertility clinic | Women with unclear infertility diagnosis                 | Endocervical | Culture     | 7%          |
| Zytoon (1994) <sup>154</sup>             | Egypt   | Cross-sectional | Convenience | 75          | W   | Infertility clinic | Women with mixed infertility diagnosis                   | Endocervical | Culture     | 65.3%       |
| Ahmadi et al (2018) <sup>49</sup>        | Iran    | Case-control    | Convenience | 165         | M   | Infertility clinic | Men with male factor infertility                         | Semen        | NAAT        | 4.2%        |
| Badami and Salari (2001) <sup>51</sup>   | Iran    | Case-control    | Convenience | 125         | W   | Infertility clinic | Women with unclear infertility diagnosis                 | Serum        | DFA         | 8.8%        |
| Badami and Salari (2001) <sup>51</sup>   | Iran    | Case-control    | Convenience | 125         | W   | Infertility clinic | Women with unclear infertility diagnosis                 | Serum        | Unclear     | 20.8%       |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | W   | Infertility clinic | Women with mixed infertility diagnosis                   | Urine        | NAAT        | 4.8%        |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | M   | Infertility clinic | 40% of men had male factor infertility                   | Urine        | NAAT        | 4.4%        |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | W   | Infertility clinic | Women with mixed infertility diagnosis                   | Serum        | EIA (IgM)   | 4%          |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | W   | Infertility clinic | Women with mixed infertility diagnosis                   | Serum        | ELISA (IgA) | 0           |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | W   | Infertility clinic | Women with mixed infertility diagnosis                   | Serum        | ELISA (IgG) | 15.6%       |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | M   | Infertility clinic | 40% of men had male factor infertility                   | Serum        | EIA (IgM)   | 1.2%        |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | M   | Infertility clinic | 40% of men had male factor infertility                   | Serum        | ELISA (IgA) | 0           |
| Dehghan et al (2017) <sup>57</sup>       | Iran    | Case-control    | Convenience | 250         | M   | Infertility clinic | 40% of men had male factor infertility                   | Serum        | ELISA (IgG) | 18%         |
| Golshani et al (2007) <sup>155</sup>     | Iran    | Cross-sectional | Convenience | 200         | M   | Infertility clinic | Majority of men had male factor infertility              | Semen        | NAAT        | 18.0%       |
| Goshayeshi et al (2015) <sup>58</sup>    | Iran    | Case-control    | Convenience | 100         | W   | Infertility clinic | Women with unclear infertility diagnosis                 | Endocervical | NAAT        | 21.0%       |
| Hajikhani et al (2013) <sup>156</sup>    | Iran    | Cross-sectional | Convenience | 51          | W   | Infertility clinic | Women with TFI   | Endocervical | Culture     | 3.9%        |
| Hajikhani et al (2013) <sup>156</sup>    | Iran    | Cross-sectional | Convenience | 51          | W   | Infertility clinic | Women with TFI   | Endocervical | NAAT        | 11.7%       |
| Joolayi et al (2017) <sup>62</sup>       | Iran    | Case-control    | Convenience | 32          | W   | Infertility clinic | Women with TFI   | Vaginal      | NAAT        | 9.4%        |
| Joolayi et al (2017) <sup>62</sup>       | Iran    | Case-control    | Convenience | 68          | W   | Infertility clinic | Women with ovarian and other infertility                 | Vaginal      | NAAT        | 2.9%        |
| Joolayi et al (2017) <sup>62</sup>       | Iran    | Case-control    | Convenience | 32          | W   | Infertility clinic | Women with TFI   | Serum        | ELISA (IgM) | 9.4%        |
| Joolayi et al (2017) <sup>62</sup>       | Iran    | Case-control    | Convenience | 68          | W   | Infertility clinic | Women with ovarian and other infertility                 | Serum        | ELISA (IgM) | 4.4%        |

(Table 2 continues on next page)

|   | Country | Study design    | Sampling*   | Sample size | Sex | Study context      | Population characteristics                      | Specimen     | Assay type      | Prevalence† |
|---|---------|-----------------|-------------|-------------|-----|--------------------|---|--------------|-----------------|-------------|
| (Continued from previous page)                                |         |                 |             |             |     |                    |   |              |                 |             |
| Joolayi et al (2017) <sup>62</sup>                            | Iran    | Case-control    | Convenience | 32          | W   | Infertility clinic | Women with TFI                                  | Serum        | ELISA (IgG)     | 0           |
| Joolayi et al (2017) <sup>62</sup>                            | Iran    | Case-control    | Convenience | 68          | W   | Infertility clinic | Women with ovarian and other infertility        | Serum        | ELISA (IgG)     | 0           |
| Kajbaf and Gholamnezhad (1998) <sup>63</sup>                  | Iran    | Case-control    | Convenience | 101         | W   | Infertility clinic | Women with mixed infertility diagnosis          | Endocervical | DFA             | 7.9%        |
| Kajbaf and Gholamnezhad (1998) <sup>63</sup>                  | Iran    | Case-control    | Convenience | 101         | W   | Infertility clinic | Women with mixed infertility diagnosis          | Serum        | ELISA (IgG)     | 17.8%       |
| Kalantar et al (2007) <sup>157</sup>                          | Iran    | Cross-sectional | Convenience | 91          | W   | Infertility clinic | Majority of women had female factor infertility | Serum        | ELISA (IgG)     | 0           |
| Kalantar et al (2007) <sup>157</sup>                          | Iran    | Cross-sectional | Convenience | 91          | W   | Infertility clinic | Majority of women had female factor infertility | Vaginal      | NAAT            | 0           |
| Kamyabi (2009) <sup>64</sup>                                  | Iran    | Case-control    | Convenience | 35          | W   | Gynaecology clinic | Women with mixed infertility diagnosis          | Serum        | ELISA (IgG)     | 22.9%       |
| Mansour Ghanaie (2014) <sup>158</sup>                         | Iran    | Cross-sectional | Convenience | 135         | W   | Infertility clinic | Majority of women without TFI                   | Endocervical | NAAT            | 19.3%       |
| Ministry of Health and Medical Education (2008) <sup>57</sup> | Iran    | Case-control    | Convenience | 46          | W   | Infertility clinic | Women with unclear infertility diagnosis        | Serum        | ELISA (IgG)     | 23.9%       |
| Marashi et al (2014) <sup>66</sup>                            | Iran    | Case-control    | Convenience | 150         | W   | Infertility clinic | Women with idiopathic infertility               | Endocervical | DFA             | 15.3%       |
| Marashi et al (2014) <sup>66</sup>                            | Iran    | Case-control    | Convenience | 150         | W   | Infertility clinic | Women with idiopathic infertility               | Endocervical | NAAT            | 32%         |
| Ministry of Health and Medical Education (2008) <sup>57</sup> | Iran    | Case-control    | Convenience | 125         | W   | Infertility clinic | Women with unclear infertility diagnosis        | Endocervical | DFA             | 8.8%        |
| Ministry of Health and Medical Education (2008) <sup>57</sup> | Iran    | Cross-sectional | Convenience | 100         | M   | Infertility clinic | Men with unclear infertility diagnosis          | Unclear      | NAAT            | 9%          |
| Moazenchi et al (2018) <sup>159</sup>                         | Iran    | Cross-sectional | Convenience | 1080        | M   | Infertility clinic | Men with unclear infertility diagnosis          | Serum        | ELISA (IgA)     | 4.3%        |
| Moazenchi et al (2018) <sup>159</sup>                         | Iran    | Cross-sectional | Convenience | 1080        | M   | Infertility clinic | Men with unclear infertility diagnosis          | Semen        | NAAT            | 10%         |
| Mousavi et al (2014) <sup>69</sup>                            | Iran    | Case-control    | Convenience | 104         | W   | Infertility clinic | Women with unclear infertility diagnosis        | Endocervical | NAAT            | 4.8%        |
| Nan Bakhsh et al (2008) <sup>160</sup>                        | Iran    | Cross-sectional | Convenience | 144         | W   | Infertility clinic | Women with mixed infertility diagnosis          | Serum        | ELISA (IgG)     | 11.1%       |
| Nikbakht et al (2008) <sup>161</sup>                          | Iran    | Case-control    | Convenience | 125         | W   | Infertility clinic | Women with TFI                                  | Unclear      | ELISA (unclear) | 23.2%       |
| Peivandi et al (2009) <sup>162</sup>                          | Iran    | Cross-sectional | Convenience | 110         | W   | Infertility clinic | Majority of women with TFI                      | Serum        | MIF (IgG)       | 24.5%       |
| Rashidi et al (2007) <sup>163</sup>                           | Iran    | Cross-sectional | Convenience | 300         | W   | Infertility clinic | Women with mixed infertility diagnosis          | Unclear      | ELISA (unclear) | 32.3%       |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience | 44          | W   | Infertility clinic | Women with TFI                                  | Urine        | NAAT            | 4.5%        |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience | 190         | W   | Infertility clinic | Women with ovarian and other infertility        | Urine        | NAAT            | 14.2%       |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience | 44          | W   | Infertility clinic | Women with TFI                                  | Serum        | ELISA (IgM)     | 2.3%        |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience | 190         | W   | Infertility clinic | Women with ovarian and other infertility        | Serum        | ELISA (IgM)     | 0.5%        |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience | 44          | W   | Infertility clinic | Women with TFI                                  | Serum        | ELISA (IgG)     | 9.1%        |
| Rashidi et al (2013) <sup>71</sup>                            | Iran    | Case-control    | Convenience | 190         | W   | Infertility clinic | Women with ovarian and other infertility        | Serum        | ELISA (IgG)     | 8.4%        |
| Sadrpour et al (2013) <sup>164</sup>                          | Iran    | Cross-sectional | Convenience | 120         | M   | Infertility clinic | Men with male factor infertility                | Semen        | NAAT            | 3%          |

(Table 2 continues on next page)

|  | Country | Study design    | Sampling*   | Sample size | Sex | Study context      | Population characteristics                  | Specimen     | Assay type     | Prevalence† |
|--|---------|-----------------|-------------|-------------|-----|--------------------|---|--------------|----------------|-------------|
| (Continued from previous page)               |         |                 |             |             |     |                    |   |              |                |             |
| Sattari et al (2017) <sup>75</sup>           | Iran    | Case-control    | Convenience | 184         | W   | Infertility clinic | Majority of women without TFI               | Serum        | ELISA (IgM)    | 5.4%        |
| Sattari et al (2017) <sup>75</sup>           | Iran    | Case-control    | Convenience | 184         | W   | Infertility clinic | Majority of women without TFI               | Serum        | ELISA (IgG)    | 35.9%       |
| Siahkali and Amini (2018) <sup>165</sup>     | Iran    | Cross-sectional | Convenience | 60          | M   | Infertility clinic | Men with idiopathic infertility             | Semen        | NAAT           | 5.0%        |
| Abid and Al-Zwaid (2015) <sup>166</sup>      | Iraq    | Case-control    | Convenience | 61          | W   | Infertility clinic | Women with mixed infertility diagnosis      | Serum        | ELISA (IgG)    | 30%         |
| Ahmed (2012) <sup>167</sup>                  | Iraq    | Case-control    | Convenience | 47          | W   | Infertility clinic | Women with unclear infertility diagnosis    | Endocervical | NAAT           | 29.8%       |
| Al-Husseinei et al (2009) <sup>84</sup>      | Iraq    | Case-control    | Convenience | 54          | W   | Infertility clinic | Women with unclear infertility diagnosis    | Endocervical | ELFA           | 9.3%        |
| Al-Husseinei et al (2009) <sup>84</sup>      | Iraq    | Case-control    | Convenience | 54          | W   | Infertility clinic | Women with unclear infertility diagnosis    | Serum        | IFAT (unclear) | 11.1%       |
| Ali and Al-Kazaz (2018) <sup>85</sup>        | Iraq    | Case-control    | Convenience | 63          | M   | Clinic             | Men with male factor infertility            | Semen        | NAAT           | 17.4%       |
| Al-Kattan and Mohammed (2013) <sup>168</sup> | Iraq    | Cross-sectional | Convenience | 54          | W   | Infertility clinic | Women with TFI or adhesions                 | Serum        | ELISA (IgG)    | 51.9%       |
| Al-Kattan and Mohammed (2013) <sup>168</sup> | Iraq    | Cross-sectional | Convenience | 67          | W   | Infertility clinic | Women without TFI or endometriosis          | Serum        | ELISA (IgG)    | 29.9%       |
| Dawood (2011) <sup>169</sup>                 | Iraq    | Cross-sectional | Convenience | 30          | W   | Hospital           | Women with unclear infertility diagnosis    | Serum        | ELISA (IgM)    | 86.6%       |
| Dawood (2011) <sup>169</sup>                 | Iraq    | Cross-sectional | Convenience | 30          | W   | Hospital           | Women with unclear infertility diagnosis    | Serum        | ELISA (IgA)    | 3.3%        |
| Dawood (2011) <sup>169</sup>                 | Iraq    | Cross-sectional | Convenience | 30          | W   | Hospital           | Women with unclear infertility diagnosis    | Serum        | ELISA (IgG)    | 53.3%       |
| Dawood (2011) <sup>169</sup>                 | Iraq    | Cross-sectional | Convenience | 100         | W   | Hospital           | Women with unclear infertility diagnosis    | Endocervical | NAAT           | 30%         |
| Ismail and Ali (2012) <sup>88</sup>          | Iraq    | Case-control    | Convenience | 52          | W   | Infertility clinic | Women with unclear infertility diagnosis    | Serum        | ELISA (IgG)    | 25%         |
| Ismail and Ali (2012) <sup>88</sup>          | Iraq    | Case-control    | Convenience | 52          | W   | Infertility clinic | Women with unclear infertility diagnosis    | Serum        | ELISA (IgM)    | 42.3%       |
| Ismail and Ali (2012) <sup>88</sup>          | Iraq    | Case-control    | Convenience | 52          | W   | Infertility clinic | Women with unclear infertility diagnosis    | Serum        | ELISA (IgA)    | 3.8%        |
| Mohammed et al (2017) <sup>90</sup>          | Iraq    | Case-control    | Convenience | 80          | W   | Gynaecology clinic | Women with mixed infertility diagnosis      | Endocervical | NAAT           | 13.8%       |
| Mohammed et al (2017) <sup>90</sup>          | Iraq    | Case-control    | Convenience | 80          | W   | Gynaecology clinic | Women with mixed infertility diagnosis      | Serum        | ELISA (IgG)    | 2.5%        |
| Yahya and Al-Siraj (2009) <sup>91</sup>      | Iraq    | Cross-sectional | Convenience | 296         | M   | Laboratory         | Men with unclear infertility diagnosis      | Serum        | Culture        | 4.0%        |
| Abusarah et al (2013) <sup>92</sup>          | Jordan  | Case-control    | Convenience | 81          | M   | Gynaecology clinic | Men with male factor infertility            | Urine        | NAAT           | 4.9%        |
| Al-Ramahi et al (2008) <sup>93</sup>         | Jordan  | Case-control    | Convenience | 66          | W   | Infertility clinic | Women with idiopathic infertility           | Endocervical | NAAT           | 3.0%        |
| Al-Ramahi et al (2008) <sup>93</sup>         | Jordan  | Case-control    | Convenience | 19          | W   | Infertility clinic | Women with TFI                              | Endocervical | NAAT           | 0           |
| Al-Ramahi et al (2008) <sup>93</sup>         | Jordan  | Case-control    | Convenience | 38          | W   | Infertility clinic | Women with male factor infertility          | Endocervical | NAAT           | 7.9%        |
| Al-Ramahi et al (2008) <sup>93</sup>         | Jordan  | Case-control    | Convenience | 29          | W   | Infertility clinic | Women with ovarian infertility              | Endocervical | NAAT           | 3.4%        |
| Al-Sweih et al (2012) <sup>100</sup>         | Kuwait  | Case-control    | Convenience | 127         | M   | Infertility clinic | Men with unclear infertility diagnosis      | Semen        | NAAT           | 3.9%        |
| Radouani et al (1998) <sup>106</sup>         | Morocco | Case-control    | Convenience | 200         | M   | Infertility clinic | Majority of men had male factor infertility | Serum        | MIF (unclear)  | 21.5%       |
| Radouani et al (1998) <sup>106</sup>         | Morocco | Case-control    | Convenience | 81          | W   | Infertility clinic | Women with unclear infertility diagnosis    | Serum        | MIF (unclear)  | 44.4%       |

(Table 2 continues on next page)



|   | Country      | Study design                 | Sampling*   | Sample size | Sex | Study context      | Population characteristics               | Specimen     | Assay type   | Prevalence† |
|---|--------------|------------------------------|-------------|-------------|-----|--------------------|--|--------------|--------------|-------------|
| (Continued from previous page)                    |              |                              |             |             |     |                    |  |              |              |             |
| Al Subhi et al (2013) <sup>170</sup>              | Oman         | Cross-sectional              | Convenience | 51          | W   | Infertility clinic | Women with TFI                           | Endocervical | EIA          | 5.9%        |
| Al Subhi et al (2013) <sup>170</sup>              | Oman         | Cross-sectional              | Convenience | 167         | W   | Infertility clinic | Women without TFI                        | Endocervical | EIA          | 4.8%        |
| Qayum and Khalid-bin-Saleem (2013) <sup>171</sup> | Pakistan     | Cross-sectional              | Convenience | 80          | W   | Gynaecology clinic | Women with unclear infertility diagnosis | Urine        | Unclear      | 7.5%        |
| Al-Hindi et al (2010) <sup>172</sup>              | Palestine    | Cross-sectional              | Convenience | 69          | W   | Infertility clinic | Women undergoing IVF in 2000             | Serum        | ELISA (IgM)  | 11.6%       |
| Al-Hindi et al (2010) <sup>172</sup>              | Palestine    | Cross-sectional              | Convenience | 268         | W   | Infertility clinic | Women undergoing IVF in 2001             | Serum        | ELISA (IgM)  | 23.9%       |
| Al-Hindi et al (2010) <sup>172</sup>              | Palestine    | Cross-sectional              | Convenience | 316         | W   | Infertility clinic | Women undergoing IVF in 2002             | Serum        | ELISA (IgM)  | 33.5%       |
| Al-Hindi et al (2010) <sup>172</sup>              | Palestine    | Cross-sectional              | Convenience | 399         | W   | Infertility clinic | Women undergoing IVF in 2003             | Serum        | ELISA (IgM)  | 9.3%        |
| Al-Hindi et al (2010) <sup>172</sup>              | Palestine    | Cross-sectional              | Convenience | 586         | W   | Infertility clinic | Women undergoing IVF in 2004             | Serum        | ELISA (IgM)  | 4.6%        |
| Al-Hindi et al (2010) <sup>172</sup>              | Palestine    | Cross-sectional              | Convenience | 316         | W   | Infertility clinic | Women undergoing IVF in 2005             | Serum        | ELISA (IgM)  | 2.8%        |
| Abdul Jabbar (1990) <sup>173</sup>                | Saudi Arabia | Cross-sectional              | Convenience | 13          | W   | Infertility clinic | Women with TFI                           | Endocervical | DFA          | 53.8%       |
| Abdul Jabbar (1990) <sup>173</sup>                | Saudi Arabia | Cross-sectional              | Convenience | 18          | W   | Infertility clinic | Women without TFI                        | Endocervical | DFA          | 11.1%       |
| Abdul Jabbar (1990) <sup>173</sup>                | Saudi Arabia | Cross-sectional              | Convenience | 34          | M   | Infertility clinic | Men with unclear infertility diagnosis   | Urethral     | DFA          | 26.4%       |
| Alfarraj et al (2015) <sup>174</sup>              | Saudi Arabia | Case-control                 | Convenience | 100         | W   | Infertility clinic | Women with mixed infertility diagnosis   | Endocervical | NAAT         | 8.0%        |
| Hossain (1988) <sup>115</sup>                     | Saudi Arabia | Cross-sectional              | Convenience | 41          | W   | Gynaecology clinic | Women with unclear infertility diagnosis | Endocervical | Culture      | 9.5%        |
| Hossain (1988) <sup>115</sup>                     | Saudi Arabia | Cross-sectional              | Convenience | 41          | W   | Gynaecology clinic | Women with unclear infertility diagnosis | Serum        | MIF (IgM)    | 0           |
| Hossain (1988) <sup>115</sup>                     | Saudi Arabia | Cross-sectional              | Convenience | 41          | W   | Gynaecology clinic | Women with unclear infertility diagnosis | Serum        | MIF (IgG)    | 16.7%       |
| Kamel (2013) <sup>116</sup>                       | Saudi Arabia | Randomised controlled trial‡ | Convenience | 640         | W   | Gynaecology clinic | Women with unclear infertility diagnosis | Endocervical | Culture      | 12.0%       |
| Kamel (2013) <sup>116</sup>                       | Saudi Arabia | Randomised controlled trial‡ | Convenience | 640         | W   | Gynaecology clinic | Women with unclear infertility diagnosis | Serum        | ELISA (IgA)  | 5%          |
| Kamel (2013) <sup>116</sup>                       | Saudi Arabia | Randomised controlled trial‡ | Convenience | 640         | W   | Gynaecology clinic | Women with unclear infertility diagnosis | Serum        | ELISA (IgG)  | 8.0%        |
| Sabra and Al-Harbi (2014) <sup>175</sup>          | Saudi Arabia | Cross-sectional              | Convenience | 148         | M   | Infertility clinic | Men with male factor infertility         | Semen        | Giemsa stain | 8.1%        |
| Almroth et al (2005) <sup>123</sup>               | Sudan        | Case-control                 | Convenience | 81          | W   | Infertility clinic | More than half of women with TFI         | Serum        | EIA (IgG)    | 14%         |
| Alkayer et al (2017) <sup>125</sup>               | Syria        | Case-control                 | Convenience | 23          | W   | Hospital           | Women with mixed infertility diagnosis   | Serum        | ELISA (IgG)  | 17.1%       |
| Gdoura et al (2001a) <sup>176</sup>               | Tunisia      | Cross-sectional              | Convenience | 92          | M   | Infertility clinic | Men with unclear infertility diagnosis   | Urethral     | NAAT         | 18.5%       |
| Gdoura et al (2001b) <sup>177</sup>               | Tunisia      | Cross-sectional              | Convenience | 92          | M   | Infertility clinic | Men with unclear infertility diagnosis   | Serum        | MIF (IgG)    | 9.8%        |
| Gdoura et al (2001a) <sup>176</sup>               | Tunisia      | Cross-sectional              | Convenience | 92          | M   | Infertility clinic | Men with unclear infertility diagnosis   | Urethral     | DFA          | 4.3%        |
| Gdoura et al (2001a) <sup>176</sup>               | Tunisia      | Cross-sectional              | Convenience | 92          | M   | Infertility clinic | Men with unclear infertility diagnosis   | Urethral     | Culture      | 1.1%        |
| Gdoura et al (2001a) <sup>176</sup>               | Tunisia      | Cross-sectional              | Convenience | 92          | M   | Infertility clinic | Men with unclear infertility diagnosis   | Urethral     | Unclear      | 8.7%        |
| Gdoura et al (2001b) <sup>177</sup>               | Tunisia      | Cross-sectional              | Convenience | 92          | W   | Infertility clinic | Partners of infertile men                | Endocervical | NAAT         | 26.1%       |

(Table 2 continues on next page)

|  | Country      | Study design    | Sampling*   | Sample size | Sex | Study context          | Population characteristics             | Specimen     | Assay type      | Prevalence† |
|--|--------------|-----------------|-------------|-------------|-----|------------------------|--|--------------|-----------------|-------------|
| (Continued from previous page)                               |              |                 |             |             |     |                        |  |              |                 |             |
| Gdoura et al (2001b) <sup>177</sup>                          | Tunisia      | Cross-sectional | Convenience | 92          | W   | Infertility clinic     | Partners of infertile men              | Serum        | MIF (IgG)       | 17.4%       |
| Gdoura et al (2008) <sup>178</sup>                           | Tunisia      | Cross-sectional | Convenience | 104         | M   | Infertility clinic     | Men with male factor infertility       | Urine        | NAAT            | 39.4%       |
| Sellami et al (2014) <sup>179</sup>                          | Tunisia      | Cross-sectional | Convenience | 85          | M   | Infertility clinic     | Men with unclear infertility diagnosis | Semen        | NAAT            | 15.2%       |
| <b>Women with miscarriage (or abortion of unknown cause)</b> |              |                 |             |             |     |                        |  |              |                 |             |
| Zaki (1989) <sup>47</sup>                                    | Egypt        | Cross-sectional | Convenience | 100         | W   | Gynaecology clinic     | Presenting with abortion               | Endocervical | Culture         | 5%          |
| Ahmadi et al (2016b) <sup>180</sup>                          | Iran         | Case-control    | Convenience | 109         | W   | Family planning clinic | Spontaneous abortion                   | Endocervical | NAAT            | 22.9%       |
| Bagheri and Roghanian (2014) <sup>181</sup>                  | Iran         | Cross-sectional | Convenience | 70          | W   | Hospital               | Recent or recurrent miscarriage        | Vaginal      | NAAT            | 1.4%        |
| Bagheri et al (2018) <sup>53</sup>                           | Iran         | Case-control    | Convenience | 97          | W   | Fertility centre       | Recent or recurrent miscarriage        | Vaginal      | NAAT            | 11.3%       |
| Bagheri et al (2018) <sup>53</sup>                           | Iran         | Case-control    | Convenience | 97          | W   | Fertility centre       | Recent or recurrent miscarriage        | Serum        | ELISA (IgA)     | 2.1%        |
| Bagheri et al (2018) <sup>53</sup>                           | Iran         | Case-control    | Convenience | 97          | W   | Fertility centre       | Recent or recurrent miscarriage        | Serum        | ELISA (IgG)     | 4.1%        |
| Jahromi et al (2010) <sup>60</sup>                           | Iran         | Case-control    | Convenience | 220         | W   | Gynaecology clinic     | Spontaneous abortion                   | Endocervical | DFA             | 25.5%       |
| Massiha et al (2010) <sup>182</sup>                          | Iran         | Cross-sectional | Convenience | 84          | W   | Hospital               | Presenting with abortion               | Unclear      | Unclear         | 2.3%        |
| Salari and Badami (2002) <sup>183</sup>                      | Iran         | Case-control    | Convenience | 125         | W   | Hospital               | Recurrent abortion                     | Endocervical | DFA             | 7.2%        |
| Sisakht et al (2017) <sup>76</sup>                           | Iran         | Case-control    | Convenience | 77          | W   | Gynaecology clinic     | Spontaneous abortion                   | Urine        | NAAT            | 9.3%        |
| Zahirnia et al (2018) <sup>78</sup>                          | Iran         | Cross-sectional | Convenience | 124         | W   | Gynaecology clinic     | Presenting with abortion               | Vaginal      | NAAT            | 15.3%       |
| Abdul-Karim et al (2009) <sup>80</sup>                       | Iraq         | Case-control    | Convenience | 79          | W   | Hospital               | Presenting with abortion               | Serum        | ELISA (IgG)     | 6.4%        |
| Abdulkhudher et al (2014) <sup>79</sup>                      | Iraq         | Case-control    | Convenience | 60          | W   | Antenatal clinic       | Recent or recurrent miscarriage        | Serum        | ELISA (IgM)     | 38.3%       |
| Abdulkhudher et al (2014) <sup>79</sup>                      | Iraq         | Case-control    | Convenience | 60          | W   | Antenatal clinic       | Recent or recurrent miscarriage        | Serum        | ELISA (IgG)     | 33.3%       |
| Ahmed (2008) <sup>82</sup>                                   | Iraq         | Case-control    | Convenience | 60          | W   | Hospital               | Recurrent miscarriage                  | Serum        | ELISA (unclear) | 0           |
| Al-Husseinei et al (2009) <sup>84</sup>                      | Iraq         | Case-control    | Convenience | 89          | W   | Family planning clinic | Recent or recurrent abortion           | Endocervical | ELFA            | 12.4%       |
| Al-Husseinei et al (2009) <sup>84</sup>                      | Iraq         | Case-control    | Convenience | 89          | W   | Family planning clinic | Recent or recurrent abortion           | Serum        | IFAT (unclear)  | 14.6%       |
| Alkhafaf (2013) <sup>86</sup>                                | Iraq         | Case-control    | Convenience | 123         | W   | Hospital               | Spontaneous abortion                   | Serum        | ELISA (IgG)     | 17.1%       |
| Al-Nuaimy and Al-Jandeel (2018) <sup>184</sup>               | Iraq         | Case-control    | Convenience | 120         | W   | Hospital               | Recent or recurrent abortion           | Endocervical | NAAT            | 17.5%       |
| Al-Nuaimy and Al-Jandeel (2018) <sup>184</sup>               | Iraq         | Case-control    | Convenience | 120         | W   | Hospital               | Recent or recurrent abortion           | Serum        | ELISA (IgG)     | 14.2%       |
| Mohammed et al (2012) <sup>89</sup>                          | Iraq         | Case-control    | Convenience | 62          | W   | Gynaecology clinic     | Three or more miscarriages             | Serum        | ELISA (IgM)     | 16.1%       |
| Mohammed et al (2012) <sup>89</sup>                          | Iraq         | Case-control    | Convenience | 34          | W   | Gynaecology clinic     | Less than three miscarriages           | Serum        | ELISA (IgM)     | 29.4%       |
| Salman (2016) <sup>185</sup>                                 | Iraq         | Cross-sectional | Convenience | 184         | W   | Gynaecology clinic     | Presenting with abortion               | Serum        | ELISA (IgM)     | 21.2%       |
| Salman (2016) <sup>185</sup>                                 | Iraq         | Cross-sectional | Convenience | 184         | W   | Gynaecology clinic     | Presenting with abortion               | Serum        | ELISA (IgG)     | 8.2%        |
| Hossain (1988) <sup>115</sup>                                | Saudi Arabia | Cross-sectional | Convenience | 12          | W   | Hospital               | Recurrent miscarriage                  | Endocervical | Culture         | 16.7%       |

(Table 2 continues on next page)

|                                     | Country      | Study design    | Sampling*   | Sample size | Sex | Study context          | Population characteristics | Specimen | Assay type  | Prevalence† |
|-------------------------------------|--------------|-----------------|-------------|-------------|-----|------------------------|----------------------------|----------|-------------|-------------|
| (Continued from previous page)      |              |                 |             |             |     |                        |                            |          |             |             |
| Hossain (1988) <sup>115</sup>       | Saudi Arabia | Cross-sectional | Convenience | 12          | W   | Hospital               | Recurrent miscarriage      | Serum    | MIF (IgM)   | 0           |
| Hossain (1988) <sup>115</sup>       | Saudi Arabia | Cross-sectional | Convenience | 12          | W   | Hospital               | Recurrent miscarriage      | Serum    | MIF (IgG)   | 16.7%       |
| <b>Women with ectopic pregnancy</b> |              |                 |             |             |     |                        |                            |          |             |             |
| Diab (1993) <sup>40</sup>           | Egypt        | Case-control    | Convenience | 30          | W   | Family planning clinic | Ectopic pregnancy          | Serum    | EIA (IgG)   | 30%         |
| Abdullah (2012) <sup>81</sup>       | Iraq         | Case-control    | Convenience | 24          | W   | Hospital               | Ectopic pregnancy          | Serum    | ELISA (IgM) | 4%          |
| Abdullah (2012) <sup>81</sup>       | Iraq         | Case-control    | Convenience | 24          | W   | Hospital               | Ectopic pregnancy          | Serum    | ELISA (IgG) | 45%         |

DFA=direct fluorescent assay. EIA=enzyme immunoassay. ELFA=enzyme-linked fluorescence assay. IFAT=indirect fluorescent antibody test. M=men or sample predominantly of men. MIF=micro-immunofluorescence. NAAT=nucleic acid amplification test. NGOs=non-governmental organisations. STI=sexually transmitted infection. TFI=tubal factor infertility. W=women or sample predominantly of women. \*Non-probability sampling refers to a sampling method in which the data collection process does not allow individuals to have equal chance of being selected; an example is convenience sampling for which individuals are selected on the basis of ease of accessibility (first-come first-served basis).<sup>122,128</sup> Probability-based sampling refers to a sampling method in which data collection process is based on a random selection of study participants; an example is random sampling from a sampling frame.<sup>129</sup> Another example of probability-based sampling is respondent-driven sampling, which is a sampling method specifically designed to sample hard-to-reach populations and is based on chain referral with the probability of selection calculated at each step in the network to produce adjusted prevalence estimates.<sup>129</sup> †The decimal places of the prevalence measures are as reported in the original report, but prevalence figures with more than one decimal place were rounded to one decimal place. ‡The extracted prevalence measure is for the baseline measurement.

**Table 2: Studies reporting *Chlamydia trachomatis* prevalence in populations at high risk, infertility clinic attendees, women with miscarriage, and women with ectopic pregnancy in the Middle East and north Africa**

High prevalence was observed in infertility clinic attendees, for both women and men (n=135), in which current infection prevalence ranged from 0 to 65.3% with a median of 9.2%, whereas ever infection prevalence ranged from 0 to 85.2% with a median of 18.6% (tables 2 and 3). Similarly, high prevalence was observed in women with miscarriage (n=27), in which current infection prevalence ranged from 1.4% to 25.5% with a median of 12.4%, whereas ever infection prevalence ranged from 4.1% to 33.3% with a median of 14.2% (tables 2 and 3).

Table 3 summarises the prevalence for other at-risk populations, and table 2 and the appendix (pp 9–14) include the full data.

The summarised and study-specific risk of bias and precision assessments are shown in the appendix (pp 15–27). Briefly, 166 (30.1%) of 552 prevalence measures were based on samples including 200 participants or more, and were classified as having higher precision. Although convenience sampling was the most common sampling methodology (495 [89.7%] of 552), probability-based sampling methods, such as respondent-driven sampling, are of increasing use for populations at high risk (18 [45%] of 40 studies in female sex workers and men who have sex with men). Almost all studies (524 [94.9%] of 552) specified the type of biological assay used for infection ascertainment (low risk of bias for this domain). Response rate was, however, unclear for 417 (75.5%) of 552 studies. Prevalence studies were overall of reasonable quality; only eight (1.4%) of 552 had high risk of bias in two or more quality domains.

Table 3 shows the meta-analyses' results for the pooled average *C trachomatis* prevalence for each at-risk population, stratified by type of assay used for infection ascertainment. Current infection prevalence was estimated at 3.0%

(95% CI 2.3–3.8) in general populations, 2.8% (1.0–5.2) in populations at intermediate risk, 13.2% (7.2–20.7) in female sex workers, 1.2% (0.2–2.8) for genital infections and 7.7% (4.2–12.0) for rectal infections in men who have sex with men, 11.3% (9.0–13.7) in infertility clinic attendees, 12.4% (7.9–17.7) in women with miscarriage, 12.4% (9.4–15.7) in symptomatic women, and 17.4% (12.5–22.8) in symptomatic men.

Meanwhile, pooled average prevalence of ever infection was estimated at 6.9% (4.3–10.0) in general populations, 1.4% (0.8–2.4) in populations at intermediate risk, 80.9% (43.8–100) in female sex workers, 21.5% (16.3–27.2) in infertility clinic attendees, 12.4% (6.6–19.5) in women with miscarriage, 37.1% (22.4–53.0) in women with ectopic pregnancy, 22.7% (15.4–31.0) in symptomatic women, and 16.9% (9.4–25.8) in symptomatic men (table 3).

Evidence for heterogeneity in *C trachomatis* prevalence estimates was observed; p values for Cochran's Q statistic was <0.0001 in most meta-analyses (table 3). Prediction intervals were generally wide affirming high heterogeneity. I<sup>2</sup> was also mostly more than 70%, indicating that most variability is due to true differences in effect size across studies rather than chance.

Figures 2 and 3 and the appendix (pp 28–40) summarise the results of subgroup meta-analyses in various subpopulations. These data show the results stratified by sex or by genital versus rectal infection (the latter only for men who have sex with men), for studies reporting current infection prevalence based on NAAT and those reporting ever infection prevalence, as well as by assay type for studies reporting current infection prevalence. Subgroup meta-analyses in infertile populations stratified by infertility diagnosis and by assay type are shown in the appendix (pp 41–42).

Table 4 summarises results of the meta-regression analyses. In the univariable analyses, at-risk population, assay type, sampling methodology, sample size, year of publication, year of data collection, country, response rate, and sex were associated with prevalence at  $p \leq 0.2$ . Alignment with meta-regression underlying assumption

|  | Studies (n) | Samples |                               | <i>C trachomatis</i> prevalence (median [range]) | Pooled average <i>C trachomatis</i> prevalence (estimate [95% CI]) | Heterogeneity measures |                           |                      |
|--|-------------|---------|-------------------------------|--|--|------------------------|---------------------------|----------------------|
|  |             | Tested  | <i>C trachomatis</i> positive |  |  | Q (p value)*           | I <sup>2</sup> † (95% CI) | Prediction interval‡ |
| General populations                        |             |         |                               |  |  |                        |                           |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                           |                      |
| NAAT                                       | 48          | 25 397  | 748                           | 2.9% (0–15.5)                                    | 3.1 (2.2–4.2)  | 714.3 (p<0.0001)       | 91.4% (89.4–93.0)         | 0.0–12.4             |
| Culture                                    | 4           | 4464    | 55                            | 5.8% (1.0–15.0)                                  | 4.3 (0.3–11.4)   | 22.5 (p<0.0001)        | 86.6% (67.7–94.5)         | 0.0–50.9             |
| Other§                                     | 23          | 128 013 | 328                           | 3.5% (0–19.9)                                    | 2.4 (1.6–3.4)  | 722.3 (p<0.0001)       | 97.0% (96.2–97.5)         | 0.0–7.2              |
| Overall current genital infection          | 75          | 157 874 | 1131                          | 3.0% (0–19.9)                                    | 3.0 (2.3–3.8)  | 2703.5 (p<0.0001)      | 97.3% (96.9–97.6)         | 0.0–10.9             |
| Anti- <i>C trachomatis</i> immunoglobulins |             |         |                               |  |  |                        |                           |                      |
| IgG (ever infection)                       | 35          | 5877    | 525                           | 4.7% (0–37.9)                                    | 6.9 (4.3–10.0)   | 226.1 (p<0.0001)       | 86.7% (82.2–90.1)         | 0.0–30.2             |
| IgM (recent infection)                     | 13          | 2843    | 74                            | 1.6% (0–14.0)                                    | 1.8 (0.3–3.9)  | 77.7 (p<0.0001)        | 84.6% (75.1–90.4)         | 0.0–12.4             |
| IgA  | 4           | 377     | 12                            | 4.3% (0–40.4)                                    | 6.2 (0.0–21.6)   | 37.8 (p<0.0001)        | 92.1% (82.9–96.3)         | 0.0–93.7             |
| Not specified (IgG, IgM, or IgA)           | 9           | 1081    | 61                            | 4.5% (0–14.8)                                    | 4.3 (1.9–7.4)  | 34.5 (p<0.0001)        | 76.8% (55.7–87.8)         | 0.0–17.3             |
| Unclear                                    | 1           | 250     | 8                             | ..   | 3.2 (1.4–6.2)  | ..                     | ..                        | ..                   |
| Populations at intermediate risk           |             |         |                               |  |  |                        |                           |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                           |                      |
| NAAT                                       | 12          | 2815    | 69                            | 1.5% (0–38.0)                                    | 2.6 (0.8–5.2)  | 117.4 (p<0.0001)       | 75.6% (56.0–86.5)         | 0.0–16.1             |
| Culture                                    | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Other§                                     | 1           | 308     | 15                            | ..   | 4.9 (2.8–7.9)  | ..                     | ..                        | ..                   |
| Overall current genital infection          | 13          | 3123    | 84                            | 2.0% (0–38.0)                                    | 2.8 (1.0–5.2)  | 127.0 (p<0.0001)       | 90.6% (85.7–93.8)         | 0.0–15.8             |
| Anti- <i>C trachomatis</i> immunoglobulins |             |         |                               |  |  |                        |                           |                      |
| IgG (ever infection)                       | 1           | 1041    | 15                            | ..   | 1.4 (0.8–2.4)  | ..                     | ..                        | ..                   |
| IgM (recent infection)                     | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| IgA  | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Not specified (IgG, IgM, or IgA)           | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Unclear                                    | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Populations at high risk                   |             |         |                               |  |  |                        |                           |                      |
| Female sex workers                         |             |         |                               |  |  |                        |                           |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                           |                      |
| NAAT                                       | 12          | 4877    | 590                           | 8.4% (0.9–72.9)                                  | 12.9 (6.5–21.0)  | 602.1 (p<0.0001)       | 98.2% (97.6–98.6)         | 0.0–52.0             |
| Culture                                    | 1           | 116     | 8                             | ..   | 6.9 (3.0–13.1)   | ..                     | ..                        | ..                   |
| Other§                                     | 1           | 30      | 9                             | ..   | 30.0 (14.7–49.4)   | ..                     | ..                        | ..                   |
| Overall current genital infection          | 14          | 5023    | 607                           | 8.4% (0.9–72.9)                                  | 13.2 (7.2–20.7)  | 611.7 (p<0.0001)       | 97.9% (97.3–98.3)         | 0.0–50.9             |
| Anti- <i>C trachomatis</i> immunoglobulins |             |         |                               |  |  |                        |                           |                      |
| IgG (ever infection)                       | 4           | 472     | 364                           | 90.0% (19.8–100)                                 | 80.9 (43.8–100.0)  | 209.9 (p<0.0001)       | 98.6% (97.7–99.1)         | 0.0–100.0            |
| IgM (recent infection)                     | 1           | 154     | 45                            | ..   | 29.2 (22.2–37.1)   | ..                     | ..                        | ..                   |
| IgA  | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Not specified (IgG, IgM, or IgA)           | 1           | 30      | 11                            | ..   | 36.7 (19.9–56.1)   | ..                     | ..                        | ..                   |
| Unclear                                    | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Men who have sex with men                  |             |         |                               |  |  |                        |                           |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                           |                      |
| NAAT                                       | 12          | 2680    | 51                            | 1.2% (0–8.8)                                     | 1.2 (0.2–2.8)  | 76.2 (p<0.0001)        | 85.6% (76.5–91.1)         | 0.0–9.5              |
| Culture                                    | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Other§                                     | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Current rectal infection                   |             |         |                               |  |  |                        |                           |                      |
| PCR  | 7           | 1506    | 129                           | 6.3% (3.6–18.3)                                  | 7.7 (4.2–12.0)   | 40.6 (p<0.0001)        | 85.2% (71.5–92.3)         | 0.0–24.9             |
| Overall current infection                  | 19          | 4186    | 180                           | 3.6% (0–18.3)                                    | 3.0 (1.2–5.4)  | 231.8 (p<0.0001)       | 92.2% (89.3–94.4)         | 0.0–17.9             |
| (Table 3 continues on next page)           |             |         |                               |  |  |                        |                           |                      |

(Table 3 continues on next page)

|  | Studies (n) | Samples |                               | <i>C trachomatis</i> prevalence (median [range]) | Pooled average <i>C trachomatis</i> prevalence (estimate [95% CI]) | Heterogeneity measures |                         |                      |
|--|-------------|---------|-------------------------------|--|--|------------------------|-------------------------|----------------------|
|  |             | Tested  | <i>C trachomatis</i> positive |  |  | Q (p value)*           | I <sup>2</sup> (95% CI) | Prediction interval‡ |
| (Continued from previous page)             |             |         |                               |  |  |                        |                         |                      |
| Anti- <i>C trachomatis</i> immunoglobulins |             |         |                               |  |  |                        |                         |                      |
| IgG (ever infection)                       | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| IgM (recent infection)                     | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| IgA  | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Not specified (IgG, IgM, or IgA)           | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Unclear                                    | 1           | 2531    | 890                           | ..   | 35.2 (33.3–37.1)   | ..                     | ..                      | ..                   |
| Infertility clinic attendees               |             |         |                               |  |  |                        |                         |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                         |                      |
| NAAT                                       | 37          | 4653    | 539                           | 8.0% (0–39.4)                                    | 10.2 (7.5–13.1)  | 310.9 (p<0.0001)       | 88.4% (85.0–91.0)       | 0.1–31.0             |
| Culture                                    | 7           | 1149    | 176                           | 9.5% (1.1–65.3)                                  | 14.4 (4.8–27.7)  | 140.8 (p<0.0001)       | 95.7% (93.3–97.3)       | 0.0–69.2             |
| Other§                                     | 20          | 1844    | 203                           | 10.2% (3.0–53.8)                                 | 12.3 (8.5–16.5)  | 112.8 (p<0.0001)       | 83.2% (75.1–88.6)       | 0.5–33.7             |
| Overall current genital infection          | 64          | 7646    | 918                           | 9.2% (0–65.3)                                    | 11.3 (9.0–13.7)  | 574.9 (p<0.0001)       | 89.0% (86.7–90.9)       | 0.2–33.3             |
| Anti- <i>C trachomatis</i> immunoglobulins |             |         |                               |  |  |                        |                         |                      |
| IgG (ever infection)                       | 37          | 3608    | 689                           | 18.6% (0–85.2)                                   | 21.5 (16.3–27.2)   | 531.8 (p<0.0001)       | 93.2% (91.6–94.6)       | 0.2–59.7             |
| IgM (recent infection)                     | 17          | 3145    | 332                           | 4.6% (0–86.7)                                    | 10.2 (5.0–16.8)  | 435.4 (p<0.0001)       | 96.3% (95.2–97.2)       | 0.0–47.0             |
| IgA  | 6           | 2302    | 82                            | 3.6% (0–5.0)                                     | 1.8 (0.1–4.7)  | 54.4 (p<0.0001)        | 90.8% (82.8–95.1)       | 0.0–16.4             |
| Not specified (IgG, IgM, or IgA)           | 5           | 760     | 211                           | 23.2% (11.1–44.4)                                | 26.1 (17.9–35.3)   | 27.5 (p<0.0001)        | 85.4% (67.8–93.4)       | 2.6–61.8             |
| Unclear                                    | 6           | 430     | 73                            | 14.8% (3.3–33.0)                                 | 14.9 (7.0–24.8)  | 30.0 (p<0.0001)        | 83.3% (65.1–92.0)       | 0.0–53.3             |
| Women with miscarriage                     |             |         |                               |  |  |                        |                         |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                         |                      |
| NAAT                                       | 6           | 597     | 87                            | 13.3% (1.4–22.9)                                 | 12.9 (7.4–19.5)  | 24.4 (p=0.0002)        | 79.5% (55.3–90.6)       | 0.1–38.9             |
| Culture                                    | 2           | 112     | 7                             | 10.9% (5.0–16.7)                                 | 7.1 (0.0–21.8)   | 2.1 (p=0.1483)         | 52.1% (0.0–88.0)        | ..                   |
| Other§                                     | 3           | 434     | 76                            | 12.4% (7.2–25.5)                                 | 14.4 (4.9–27.6)  | 21.9 (p<0.0001)        | 90.8% (76.1–96.5)       | 0.0–100.0            |
| Overall current genital infection          | 11          | 1143    | 170                           | 12.4% (1.4–25.5)                                 | 12.4 (7.9–17.7)  | 58.0 (p<0.0001)        | 82.8% (70.5–89.9)       | 0.6–34.3             |
| Anti- <i>C trachomatis</i> immunoglobulins |             |         |                               |  |  |                        |                         |                      |
| IgG (ever infection)                       | 7           | 675     | 84                            | 14.2% (4.1–33.3)                                 | 12.4 (6.6–19.5)  | 33.5 (p<0.0001)        | 82.1% (64.2–91.0)       | 0.0–39.6             |
| IgM (recent infection)                     | 5           | 352     | 82                            | 21.2% (0–38.3)                                   | 21.2 (11.9–32.2)   | 16.2 (p=0.0028)        | 75.3% (39.1–89.9)       | 0.0–61.8             |
| IgA  | 1           | 97      | 2                             | ..   | 2.1 (0.3–7.3)  | ..                     | ..                      | ..                   |
| Not specified (IgG, IgM, or IgA)           | 2           | 149     | 13                            | 7.3% (0–14.6)                                    | 4.7 (0.0–27.9)   | 16.1 (p<0.0001)        | 93.8% (80.0–98.1)       | ..                   |
| Unclear                                    | 1           | 84      | 2                             | ..   | 2.3 (0.3–8.3)  | ..                     | ..                      | ..                   |
| Women with ectopic pregnancy               |             |         |                               |  |  |                        |                         |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                         |                      |
| NAAT                                       | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Culture                                    | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Other§                                     | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Overall current genital infection          | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Anti- <i>C trachomatis</i> immunoglobulins |             |         |                               |  |  |                        |                         |                      |
| IgG (ever infection)                       | 2           | 54      | 20                            | 37.5% (30.0–45.0)                                | 37.1 (22.4–53.0)   | 1.4 (p=0.2418)         | 27.0%                   | ..                   |
| IgM (recent infection)                     | 1           | 24      | 1                             | ..   | 4.2 (0.1–21.1)   | ..                     | ..                      | ..                   |
| IgA  | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Not specified (IgG, IgM, or IgA)           | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Unclear                                    | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                      | ..                   |
| Symptomatic women                          |             |         |                               |  |  |                        |                         |                      |
| Current genital infection                  |             |         |                               |  |  |                        |                         |                      |
| NAAT                                       | 49          | 14 398  | 1123                          | 8.0% (0–68.0)                                    | 8.8 (6.2–11.7)   | 1506.7 (p<0.0001)      | 96.8% (96.3–97.3)       | 0.0–35.4             |
| Culture                                    | 10          | 2951    | 752                           | 12.9% (0.7–69.4)                                 | 18.9 (4.1–40.9)  | 1511.1 (p<0.0001)      | 99.4% (99.3–99.5)       | 0.0–97.2             |
| Other§                                     | 31          | 4796    | 729                           | 14.7% (0–89.3)                                   | 16.8 (11.6–22.7)   | 723.8 (p<0.0001)       | 95.9% (94.9–96.6)       | 0.0–55.4             |

(Table 3 continues on next page)

(Table 3 continues on next page)

|  | Studies (n) | Samples |                               | <i>C trachomatis</i> prevalence (median [range]) | Pooled average <i>C trachomatis</i> prevalence (estimate [95% CI]) | Heterogeneity measures |                           |                      |
|--|-------------|---------|-------------------------------|--|--|------------------------|---------------------------|----------------------|
|  |             | Tested  | <i>C trachomatis</i> positive |  |  | Q (p value)*           | I <sup>2</sup> † (95% CI) | Prediction interval‡ |
| (Continued from previous page)   |             |         |                               |  |  |                        |                           |                      |
| Overall current genital infection  | 90          | 22 145  | 2604                          | 11.7% (0–89.3)                                   | 12.4 (9.4–15.7)  | 4323.7 (p<0.0001)      | 97.9% (97.7–98.1)         | 0.0–52.3             |
| Anti- <i>C trachomatis</i> immunoglobulins   |             |         |                               |  |  |                        |                           |                      |
| IgG (ever infection)   | 23          | 2377    | 609                           | 18.2% (2.7–86.0)                                 | 22.7 (15.4–31.0)   | 454.1 (p<0.0001)       | 95.2% (93.8–96.2)         | 0.0–67.9             |
| IgM (recent infection)   | 9           | 1042    | 160                           | 3.1% (0–86.0)                                    | 13.9 (1.9–33.6)  | 452.7 (p<0.0001)       | 98.2% (97.6–98.7)         | 0.0–91.3             |
| IgA  | 5           | 365     | 93                            | 12.5% (3.7–59.6)                                 | 24.9 (6.6–49.6)  | 98.9 (p<0.0001)        | 96.0% (93.0–97.7)         | 0.0–100.0            |
| Not specified (IgG, IgM, or IgA)   | 7           | 2530    | 761                           | 15.0% (11.6–46.6)                                | 23.5 (11.1–38.7)   | 346.8 (p<0.0001)       | 98.3% (97.6–98.4)         | 0.0–80.0             |
| Unclear  | 6           | 943     | 83                            | 7.7% (1.7–30.0)                                  | 8.7 (4.3–14.4)   | 32.2 (p<0.0001)        | 84.5% (67.9–92.5)         | 0.0–32.6             |
| Symptomatic men  |             |         |                               |  |  |                        |                           |                      |
| Current genital infection  |             |         |                               |  |  |                        |                           |                      |
| NAAT   | 14          | 7160    | 726                           | 12.2% (4.2–33.3)                                 | 13.9 (8.3–20.6)  | 488.7 (p<0.0001)       | 97.3% (96.5–98.0)         | 0.0–46.4             |
| Culture  | 5           | 4744    | 75                            | 9.3% (0.4–19.6)                                  | 8.7 (1.1–21.7)   | 147.5 (p<0.0001)       | 97.3% (95.6–98.3)         | 0.0–72.7             |
| Other§   | 13          | 2499    | 355                           | 27.6% (4.7–52.0)                                 | 26.3 (15.3–39.1)   | 351.7 (p<0.0001)       | 96.6% (95.4–97.5)         | 0.0–78.9             |
| Overall current genital infection  | 32          | 14 403  | 1156                          | 15.5% (0.4–52.0)                                 | 17.4 (12.5–22.8)   | 1628.1 (p<0.0001)      | 98.1% (97.8–98.4)         | 0.0–53.3             |
| Anti- <i>C trachomatis</i> immunoglobulins   |             |         |                               |  |  |                        |                           |                      |
| IgG (ever infection)   | 8           | 831     | 164                           | 14.4% (5.1–46.0)                                 | 16.9 (9.4–25.8)  | 69.6 (p<0.0001)        | 89.9% (82.6–94.2)         | 0.0–52.8             |
| IgM (recent infection)   | 3           | 330     | 24                            | 3.9% (2.8–12.2)                                  | 6.0 (1.4–13.0)   | 9.3 (p=0.0095)         | 78.5% (31.0–93.3)         | 0.0–100.0            |
| IgA  | ..          | ..      | ..                            | ..   | ..   | ..                     | ..                        | ..                   |
| Not specified (IgG, IgM, or IgA)   | 5           | 1596    | 687                           | 46.0% (10.0–49.1)                                | 35.6 (18.5–54.9)   | 143.3 (p<0.0001)       | 97.2% (95.4–98.3)         | 0.0–97.3             |
| Unclear  | 8           | 4876    | 233                           | 14.3% (1.6–76.9)                                 | 24.0 (9.2–42.8)  | 498.1 (p<0.0001)       | 98.6% (98.1–99.0)         | 0.0–90.3             |
| A minimum of two studies was necessary to do a meta-analysis. The same population might have contributed different measures for both current and ever infection with <i>C trachomatis</i> . NAAT=nucleic acid amplification test. *Cochran's Q statistic is a measure assessing the existence of heterogeneity in effect size of <i>C trachomatis</i> prevalence across studies. †I <sup>2</sup> is a measure assessing the magnitude of between-study variation that is due to differences in effect size of <i>C trachomatis</i> prevalence across studies rather than chance. ‡Prediction interval is a measure estimating the 95% CI of the distribution of true effect sizes of <i>C trachomatis</i> prevalence measures. §Other assays detecting current infection such as direct fluorescence assays, Giemsa staining, and enzyme-linked immunoassays applied to genital samples. |             |         |                               |  |  |                        |                           |                      |
| Table 3: Results of meta-analyses on studies reporting <i>Chlamydia trachomatis</i> prevalence in the Middle East and north Africa stratified by at-risk population and <i>C trachomatis</i> ascertainment   |             |         |                               |  |  |                        |                           |                      |

**Table 3: Results of meta-analyses on studies reporting *Chlamydia trachomatis* prevalence in the Middle East and north Africa stratified by at-risk population and *C trachomatis* ascertainment**

of normal random effects was confirmed through normal probability plots (appendix p 43).<sup>186</sup> Graphical illustrations of the fitted regression line for year of publication and year of data collection are shown in the appendix (p 44). Only at-risk population, assay type, sample size, country, and sex remained associated with *C trachomatis* prevalence in a multivariable model. No evidence was found for a temporal variation in prevalence ( $p=0.281$  for year of publication), for sampling methodology ( $p=0.347$ ), or for response rate ( $p=0.237$ ). This model explained 29.0% of prevalence variation.

Relative to general populations, the adjusted odds ratio (aOR) was 11.28 (95% CI 5.78–22.01) for female sex workers, 7.17 (4.05–12.68) for symptomatic men, 4.93 (1.03–23.52) for women with ectopic pregnancy, 4.16 (1.72–10.08) for men who have sex with men, 3.39 (2.41–4.77) for infertility clinic attendees, 3.47 (2.47–4.87) for symptomatic women, 2.78 (1.57–4.93) for women with miscarriage, and 1.81 (0.79–4.13) for populations at intermediate risk. Other factors associated with *C trachomatis* prevalence were women versus men (aOR 1.61, 95% CI 1.05–2.46), Pakistan versus other Middle East or north African countries (0.39, 0.22–0.69), ever infection (anti-*C trachomatis* IgG; 2.17, 1.54–3.06) and current infection prevalence using assays other than

NAAT or culture versus NAAT (1.47, 1.02–2.13), and studies with higher ( $\geq 200$  participants) versus lower precision (0.63, 0.48–0.83).

## Discussion

We provided a comprehensive assessment of *C trachomatis* epidemiology in the Middle East and north Africa.<sup>2,3</sup> Unexpectedly, given this region's sexually conservative norms and low observed levels of several viral STIs,<sup>10,11,187–189</sup> *C trachomatis* current infection prevalence was 3% in the population at large, similar to WHO prevalence estimates for this region of about 3% in 2012<sup>3</sup> and about 3.5% in 2016.<sup>190</sup> The prevalence was also in line with WHO estimates for the Western Pacific region (about 4%) and European region (about 3%),<sup>190</sup> where broad *C trachomatis* control programmes, including opportunistic testing, are standard in some high-income countries,<sup>191–193</sup> but higher than that for South-east Asia region (about 1.5%) and lower than that for the African region (about 5%) and the region of the Americas (about 5.5%).<sup>190</sup> This high prevalence suggests substantial infection and disease burden that needs to be tackled through sexual health and STI-specific programmes, for both women and men. Although these findings were based on a volume of epidemiological evidence, most studies used convenience




sampling (about 90%) or had unclear response rate (>75%). Meta-regression, however, did not identify an effect for these factors on observed prevalence. A summary of this study and its results in Arabic language can be found in the appendix (p 4).

Although infection prevalence in the population at large suggests active transmission networks for *C trachomatis* and other STIs, it might not necessarily reflect prevalent sexual risk behaviours. This outcome might reflect, at least in part, poor access to and utilisation of STI services—there is very limited capacity in the Middle East and north Africa for STI prevention and treatment, not to mention *C trachomatis* screening and broader sexual health programmes. As observed elsewhere, such as in Alaskan Eskimo populations<sup>194</sup> and populations in South Pacific Islands,<sup>195</sup> poor *C trachomatis* diagnosis and specific treatment can result in unusually high prevalence,<sup>194,196</sup> probably because *C trachomatis* is largely asymptomatic,<sup>4</sup> and if untreated, shedding can persist even for years,<sup>197</sup> thereby increasing the potential for reinfection within couples<sup>198</sup> and for transmission in the population.


The high prevalence found in populations at high risk such as female sex workers, in context of evidence suggesting strong partial immunity against reinfection,<sup>199</sup> is consistent with the important role of commercial sex networks in infection transmission. Independent evidence supports existence of hidden pockets of high sexual-risk behaviour driving STI incidence in the Middle East and north Africa.<sup>10,11</sup> Among male STI patients, 77% in Kuwait<sup>200</sup> and 80% in Somalia<sup>201</sup> reported paying a female sex worker for sex, and among migrant workers in Pakistan 22% reported sex with a female sex worker.<sup>202</sup> Higher levels of sexual-risk behaviour and emerging HIV epidemics have been also documented among men who have sex with men, male sex workers, and male-to-female transgenders in systematic reviews.<sup>17,203</sup> Sexual networks, however, remain poorly investigated in the Middle East and north Africa, owing to cultural sensitivities.

The possible role of *C trachomatis* infection in poor reproductive health outcomes remains unappreciated and neglected by the public health establishment in the Middle East and north Africa, despite substantial social and economic implications for women and their families.<sup>204,205</sup> A main finding of this study is the high current *C trachomatis* infection prevalence in infertility clinic attendees, with odds of infection three-times higher than in the general population. By contrast, studies among infertility clinic attendees in Europe usually show that current *C trachomatis* infection is uncommon, but serological evidence of past infection, assumed to have resulted in fallopian tube scarring, is common.<sup>206–209</sup> This finding suggests a role for *C trachomatis* in infertility in the Middle East and north Africa. Indeed, this region appears to have the highest rate of primary infertility worldwide, which remains unexplained.<sup>12</sup> The Middle East and north Africa is also a region where infertility has multiple



**Figure 2: Meta-analysis of studies reporting *Chlamydia trachomatis* current infection prevalence assessed using nucleic acid amplification test in the general population in the Middle East and North Africa**  
Data are stratified by sex. Error bars are 95% CIs.

detrimental sociocultural consequences,<sup>210</sup> and where several countries have had rapidly declining fertility rates to even below replacement level.<sup>211,212</sup> The prevalence of current *C trachomatis* infection was also high in women



**Figure 3: Meta-analysis of studies reporting *Chlamydia trachomatis* current infection prevalence assessed using nucleic acid amplification test in infertility clinic attendees in the Middle East and north Africa**  
Data are stratified by sex. Error bars are 95% CIs.

with miscarriage and in pregnant women—similar to that found in pregnant women in low-income and middle-income countries elsewhere.<sup>213–217</sup> This stigmatised and largely asymptomatic infection might not be visible to the public eye, but its reproductive health sequelae are visible, even if not explicitly linked to the underlying cause.

*C trachomatis* prevalence in women was higher than in men (two-times higher odds). This difference possibly reflects a longer duration of infection for women, considering that infection in men is more symptomatic<sup>218</sup> (nearly two-times higher prevalence in symptomatic men than in symptomatic women), and therefore more likely to be treated. Ever infection (anti-*C trachomatis* IgG)

prevalence was two-times higher than current infection prevalence, but the epidemiological relevance of ever infection prevalence might be limited given challenges in *C trachomatis* serology interpretation.<sup>219</sup>

The Middle East and north Africa is burdened by *C trachomatis* infection, but the public health response remains rudimentary and far from achieving WHO's Global Health Sector Strategy on STIs.<sup>8</sup> Evidence for some differences in *C trachomatis* prevalence by country has been reported, but remarkably, no evidence was found for a variation in prevalence over time (1982–2018). Lingering STI stigma prevents those infected from accessing proper health care, including those most at risk. The role of screening and treatment for asymptomatic *C trachomatis* within established programmes, such as for family planning, primary health care, or HIV, needs careful consideration given the cost and uncertain effect on prevalence at modest levels of uptake.<sup>220</sup>

Current STI surveillance focused on inefficient routine case reporting is not capturing the reality of the transmission dynamics.<sup>221</sup> Although routine case reporting could be improved with more consistency and universality in reporting and emphasis on aetiological approaches,<sup>221</sup> its usefulness for a robust long-term evaluation of infection trends is rather limited. Sentinel surveillance of different at-risk populations should be explored, as recommended by the WHO Global Health Sector Strategy on STIs,<sup>8</sup> to better identify outbreaks or emerging epidemics, strategically direct resources for prevention, treatment, and control, and monitor and evaluate STI programmes.<sup>1</sup> The recent progress in HIV surveillance in the Middle East and north Africa, in the form of repeated rounds of HIV-integrated biobehavioural surveillance surveys,<sup>222,223</sup> should be extended to STIs.<sup>221,224</sup>

Our study has important but unavoidable limitations. Quantity and quality of available data varied by country and population, particularly for populations at high risk where most data came from only a few countries—eg, most studies of men who have sex with men were from Pakistan. No data were identified for Afghanistan, Bahrain, Libya, and Yemen. Prevalence levels might not have been strictly representative and might have been affected by publication bias, as suggested by the small-study effect observed. Studies in women with miscarriage might have included women with induced abortions; however, these were not explicitly indicated, possibly for legal reasons, as abortion is illegal in most of the Middle East and north Africa.<sup>225,226</sup> The wide array of diagnostics used for ascertainment might have also introduced detection bias.

Factors that might have contributed to differences in *C trachomatis* positivity rates across studies include sampling variation and potential selection bias, spatial or temporal variability in prevalence, and possibly unreported underlying comorbidities. This study did not assess other STIs that might have also contributed to infertility, pregnancy-related morbidity, and other health conditions in women with *C trachomatis* infection. Such

potential biases might have contributed to some of the unexplained heterogeneity observed in the prevalence levels. Given potential limitations in the representativeness of the prevalence measures as well as heterogeneity across studies, the calculated pooled prevalence should be interpreted as a pooled average, rather than strictly

|  | Studies (n) | Samples (n) | Univariable analyses |         |                  | Variance explained R <sup>2</sup> | Multivariable analysis |         |                  |
|--|-------------|-------------|----------------------|---------|------------------|-----------------------------------|------------------------|---------|------------------|
|  |             |             | OR (95% CI)          | p value | LR test p value* |                                   | Adjusted OR (95% CI)   | p value | LR test p value† |
| At-risk population                           |             |             |                      |         |                  |                                   |                        |         |                  |
| General populations                          | 137         | 168 302     | 1.00 (ref)           | ..      | <0.0001          | 19.0%                             | 1.00 (ref)             | ..      | <0.0001          |
| Populations at intermediate risk             | 14          | 4164        | 0.70 (0.32–1.54)     | 0.374   | ..               | ..                                | 1.81 (0.79–4.13)       | 0.157   | ..               |
| Female sex workers                           | 20          | 5679        | 8.99 (4.57–17.71)    | <0.0001 | ..               | ..                                | 11.28 (5.78–22.01)     | <0.0001 | ..               |
| Men who have sex with men                    | 20          | 6717        | 0.83 (0.42–1.64)     | 0.591   | ..               | ..                                | 4.16 (1.72–10.08)      | 0.002   | ..               |
| Infertility clinic attendees                 | 135         | 17 891      | 3.77 (2.67–5.31)     | <0.0001 | ..               | ..                                | 3.39 (2.41–4.77)       | <0.0001 | ..               |
| Women with miscarriage                       | 27          | 2500        | 3.53 (1.94–6.40)     | <0.0001 | ..               | ..                                | 2.78 (1.57–4.93)       | 0.001   | ..               |
| Women with ectopic pregnancy                 | 3           | 78          | 8.25 (1.58–43.08)    | 0.012   | ..               | ..                                | 4.93 (1.03–23.52)      | 0.045   | ..               |
| Symptomatic women                            | 140         | 29 402      | 3.74 (2.66–5.26)     | <0.0001 | ..               | ..                                | 3.47 (2.47–4.87)       | <0.0001 | ..               |
| Symptomatic men                              | 56          | 22 036      | 5.76 (3.68–9.03)     | <0.0001 | ..               | ..                                | 7.17 (4.05–12.68)      | <0.0001 | ..               |
| Assay type                                   |             |             |                      |         |                  |                                   |                        |         |                  |
| NAAT (current infection)                     | 197         | 64 083      | 1.00 (ref)           | ..      | <0.0001          | 7.1%                              | 1.00 (ref)             | ..      | <0.0001          |
| Culture (current infection)                  | 29          | 13 536      | 1.92 (1.05–3.50)     | 0.034   | ..               | ..                                | 1.10 (0.62–1.95)       | 0.742   | ..               |
| Other (current infection)‡                   | 92          | 137 924     | 1.90 (1.30–2.79)     | 0.001   | ..               | ..                                | 1.47 (1.02–2.13)       | 0.041   | ..               |
| Anti- <i>C. trachomatis</i> immunoglobulins§ |             |             |                      |         |                  |                                   |                        |         |                  |
| IgG (ever infection)                         | 117         | 14 935      | 2.99 (2.10–4.26)     | <0.0001 | ..               | ..                                | 2.17 (1.54–3.06)       | <0.0001 | ..               |
| IgM (recent infection)                       | 49          | 7890        | 1.17 (0.72–1.90)     | 0.517   | ..               | ..                                | 0.90 (0.57–1.40)       | 0.627   | ..               |
| IgA  | 16          | 3141        | 0.92 (0.42–2.02)     | 0.836   | ..               | ..                                | 0.78 (0.39–1.56)       | 0.481   | ..               |
| Not specified (IgG, IgM, or IgA)             | 29          | 6146        | 2.81 (1.54–5.13)     | 0.001   | ..               | ..                                | 2.25 (1.28–3.97)       | 0.005   | ..               |
| Unclear                                      | 23          | 9114        | 2.53 (1.30–4.94)     | 0.007   | ..               | ..                                | 1.49 (0.81–2.75)       | 0.200   | ..               |
| Sampling methodology¶                        |             |             |                      |         |                  |                                   |                        |         |                  |
| Non-probability-based sampling               | 495         | 227 208     | 1.00 (ref)           | ..      | <0.0001          | 3.5%                              | 1.00 (ref)             | ..      | 0.347            |
| Probability-based sampling                   | 57          | 29 561      | 0.37 (0.24–0.56)     | <0.0001 | ..               | ..                                | 0.80 (0.50–1.27)       | 0.347   | ..               |
| Sample size                                  |             |             |                      |         |                  |                                   |                        |         |                  |
| <200   | 386         | 32 782      | 1.00 (ref)           | ..      | <0.0001          | 6.0%                              | 1.00 (ref)             | ..      | 0.001            |
| ≥200   | 166         | 223 987     | 0.42 (0.32–0.56)     | <0.0001 | ..               | ..                                | 0.63 (0.48–0.83)       | 0.001   | ..               |
| Response rate                                |             |             |                      |         |                  |                                   |                        |         |                  |
| ≥80%   | 112         | 38 732      | 1.00 (ref)           | ..      | 0.187            | 0.1%                              | 1.00 (ref)             | ..      | 0.237            |
| <80% or unclear                              | 440         | 218 037     | 0.80 (0.57–1.12)     | 0.187   | ..               | ..                                | 0.83 (0.61–1.13)       | 0.237   | ..               |
| Year of publication                          | 552         | 256 769     | 0.96 (0.95–0.98)     | <0.0001 | <0.0001          | 4.4%                              | 0.99 (0.98–1.01)       | 0.281   | 0.281            |
| Year of data collection                      | 552         | 256 769     | 0.96 (0.95–0.98)     | <0.0001 | <0.0001          | 4.2%                              | ..                     | ..      | ..               |
| Country                                      |             |             |                      |         |                  |                                   |                        |         |                  |
| Other Middle East or north African countries | 245         | 189 529     | 1.00 (ref)           | ..      | <0.0001          | 5.2%                              | 1.00 (ref)             | ..      | 0.013            |
| Egypt  | 89          | 7434        | 1.58 (1.08–2.31)     | 0.018   | ..               | ..                                | 1.05 (0.73–1.51)       | 0.774   | ..               |
| Iran   | 176         | 38 647      | 0.80 (0.59–1.08)     | 0.145   | ..               | ..                                | 0.90 (0.68–1.19)       | 0.472   | ..               |
| Pakistan                                     | 42          | 21 159      | 0.31 (0.19–0.52)     | <0.0001 | ..               | ..                                | 0.39 (0.22–0.69)       | 0.002   | ..               |
| Sex  |             |             |                      |         |                  |                                   |                        |         |                  |
| Men  | 133         | 42 393      | 1.00 (ref)           | ..      | 0.131            | 0.2%                              | 1.00 (ref)             | ..      | 0.029            |
| Women  | 419         | 214 376     | 1.27 (0.93–1.74)     | 0.131   | ..               | ..                                | 1.61 (1.05–2.46)       | 0.029   | ..               |

Adjusted R<sup>2</sup> in the final multivariable model was 29.0%. LR=likelihood ratio. NAAT=nucleic acid amplification test. OR=Odds ratio. \*Predictors with p≤0.2 in the univariable model were considered significant. †Predictors with p≤0.05 in the multivariable model were considered significant. ‡Other assays detecting current infection such as direct fluorescence assays, Giemsa staining, and enzyme-linked immunoassays applied to genital samples. §Includes assays such as enzyme-linked immunoassay and micro-immunofluorescence. ¶Non-probability sampling refers to a sampling method in which the data collection process does not allow individuals to have equal chance of being selected; an example is convenience sampling for which individuals are selected on the basis of ease of accessibility (first-come first-served basis).<sup>127,128</sup> Probability-based sampling refers to a sampling method in which data collection process is based on a random selection of study participants; an example is random sampling from a sampling frame.<sup>128</sup> Another example of probability-based sampling is respondent-driven sampling, which is a sampling method specifically designed to sample hard-to-reach populations and is based on chain referral with the probability of selection calculated at each step in the network to produce adjusted prevalence estimates.<sup>129</sup> ||Only year of publication was considered for the multivariable meta-regression analysis because of collinearity with year of data collection.

**Table 4: Results of meta-regressions to identify associations and sources of between-study heterogeneity for *Chlamydia trachomatis* prevalence in the Middle East and north Africa**

an estimate of the mean prevalence in the considered population or subpopulation.

In conclusion, *C trachomatis* current infection prevalence in the population at large in the Middle East and north Africa is at 3%, similar to other regions, but higher than expected given these countries' sexually conservative norms. The high prevalence (>10%) in infertility clinic attendees and in women with miscarriage, provides suggestive evidence for the potential role of *C trachomatis* in poor reproductive outcomes in the Middle East and north Africa. In the context of very limited programming for sexual health and STIs, our findings highlight an important, yet neglected and poorly recognised infection and disease burden, despite the social and economic impact. There is an urgent need for targeted and culturally appropriate programmes promoting sexual health for different at-risk populations. Tackling this infection with appropriate interventions is essential to control disease sequelae, to address the WHO Global Health Sector Strategy on STIs,<sup>8</sup> and to accomplish key health Sustainable Development Goals.

#### Contributors

AS contributed to the study design, did the systematic searches of the literature, selection of studies for inclusion, and the data extraction and data analyses. HC contributed to the study design, double extracted the data, updated the systematic review, and did the data analyses. AS and HC wrote the first draft of the paper. JGH contributed to identification of unpublished data. NL contributed to the data extraction, analyses, and drafting of the Article. LJA-R conceived and led the design of the study, data extraction, data analyses, and drafting of the Article. All authors contributed to discussion and interpretation of the results and to the writing of the manuscript. All authors have read and approved the final manuscript.

#### Declaration of interests

We declare no competing interests.

#### Acknowledgments

This study was funded by the National Priorities Research Program (grant number 9-040-3-008) from the Qatar National Research Fund (a member of Qatar Foundation). Infrastructure support was provided by the Biostatistics, Epidemiology, and Biomathematics Research Core at Weill Cornell Medicine-Qatar, Cornell University, Qatar Foundation—Education City, Doha, Qatar. The statements made herein are solely the responsibility of the authors. The publication of this Article was funded by the Qatar National Library. We thank Adona Canlas and Ashwini Deshmukh for their assistance with locating full-text articles, and Ghina Mumtaz and Vajihah Akbarzadeh for their assistance with translating manuscripts from foreign languages.

#### References

- WHO. Baseline report on global sexually transmitted infection surveillance 2012. Geneva: World Health Organization, 2013. <http://www.who.int/reproductivehealth/publications/rtis/9789241505895/en/> (accessed March 9, 2014).
- Holmes KK. Sexually transmitted diseases, 4th edn. New York: McGraw-Hill Medical, 2008.
- Newman L, Rowley J, Vander Hoorn S, et al. Global estimates of the prevalence and incidence of four curable sexually transmitted infections in 2012 based on systematic review and global reporting. *PLoS One* 2015; **10**: e0143304.
- WHO. Global prevalence and incidence of selected curable sexually transmitted infections: overview and estimates. Geneva: World Health Organization, 2001.
- Mishori R, McClaskey EL, WinklerPrins VJ. *Chlamydia trachomatis* infections: screening, diagnosis, and management. *Am Fam Physician* 2012; **86**: 1127–32.
- Rekart ML, Gilbert M, Meza R, et al. Chlamydia public health programs and the epidemiology of pelvic inflammatory disease and ectopic pregnancy. *J Infect Dis* 2013; **207**: 30–38.
- UN General Assembly. Transforming our world: the 2030 Agenda for Sustainable Development. 2015. <https://sustainabledevelopment.un.org/content/documents/21252030%20Agenda%20for%20Sustainable%20Development%20web.pdf> (accessed June 6, 2017).
- WHO. Global health sector strategy on sexually transmitted infections, 2016–2021. Geneva: World Health Organization, 2016.
- Abu-Raddad LJ, Ghanem KG, Feizzadeh A, Setayesh H, Calleja JM, Riedner G. HIV and other sexually transmitted infection research in the Middle East and north Africa: promising progress? *Sex Transm Infect* 2013; **89** (suppl 3): iii1–4.
- Abu-Raddad LJ, Hilmi N, Mumtaz G, et al. Epidemiology of HIV infection in the Middle East and north Africa. *AIDS* 2010; **24** (suppl 2): S5–23.
- Abu-Raddad LJ, Akala FA, Semini I, Riedner G, Wilson D, Tawil O. Characterizing the HIV/AIDS epidemic in the Middle East and north Africa: time for strategic action. Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project. Washington DC: The World Bank Press, 2010.
- Mascarenhas MN, Flaxman SR, Boerma T, Vanderpoel S, Stevens GA. National, regional, and global trends in infertility prevalence since 1990: a systematic analysis of 277 health surveys. *PLoS Med* 2012; **9**: e1001356.
- Higgins JPT, Green S, Cochrane Collaboration. Cochrane handbook for systematic reviews of interventions. Chichester: Wiley-Blackwell, 2008.
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097.
- International AIDS Society. Abstract archives of international aids society conferences. 2016. <http://www.abstract-archive.org/> (accessed July 17, 2017).
- Abu-Raddad LJ, World Bank. Characterizing the HIV/AIDS epidemic in the Middle East and north Africa: time for strategic action. Washington DC: World Bank, 2010.
- Mumtaz G, Hilmi N, McFarland W, et al. Are HIV epidemics among men who have sex with men emerging in the Middle East and north Africa? A systematic review and data synthesis. *PLoS Med* 2010; **8**: e1000444.
- Mumtaz GR, Weiss HA, Thomas SL, et al. HIV among people who inject drugs in the Middle East and north Africa: systematic review and data synthesis. *PLoS Med* 2014; **11**: e1001663.
- Harfouche M, Chemaitelly H, Mahmud S, et al. Epidemiology of hepatitis C virus among hemodialysis patients in the Middle East and north Africa: systematic syntheses, meta-analyses, and meta-regressions. *Epidemiol Infect* 2017; **145**: 3243–63.
- Mahmud S, Al-Kanaani Z, Chemaitelly H, Chaabna K, Kouyoumjian SP, Abu-Raddad LJ. Hepatitis C virus genotypes in the Middle East and north Africa: distribution, diversity, and patterns. *J Med Virol* 2018; **90**: 131–41.
- Harfouche M, Chemaitelly H, Kouyoumjian SP, et al. Hepatitis C virus viremic rate in the Middle East and north Africa: systematic synthesis, meta-analyses, and meta-regressions. *PLoS One* 2017; **12**: e0187177.
- Chemaitelly H, Mahmud S, Chaabna K, Kouyoumjian SP, Mumtaz GR, Abu-Raddad LJ. The epidemiology of hepatitis C virus in the World Health Organization Eastern Mediterranean region: implications for strategic action. 2019. World Health Organization Report (in press).
- El-Kettani A, Mahiane G, Bennani A, et al. Trends in adult chlamydia and gonorrhea prevalence, incidence and urethral discharge case reporting in Morocco over 1995–2015: estimates using the spectrum-sexually transmitted infection model. *Sex Transm Infect* 2017; **44**: 557–64.
- Smolak A, Rowley J, Nagelkerke N, et al. Trends and predictors of syphilis prevalence in the general population: global pooled analyses of 1103 prevalence measures including 136 million syphilis tests. *Clin Infect Dis* 2018; **66**: 1184–91.
- Korenromp EL, Mahiane G, Rowley J, et al. Estimating prevalence trends in adult gonorrhoea and syphilis in low- and middle-income countries with the spectrum-STI model: results for Zimbabwe and Morocco from 1995 to 2016. *Sex Transm Infect* 2017; **93**: 599–606.



- 26 Freeman MF, Tukey JW. Transformations related to the angular and the square root. *Ann Math Stat* 1950; **21**: 607–11.
- 27 Miller JJ. The inverse of the Freeman–Tukey double arcsine transformation. *Am Stat* 1978; **32**: 138.
- 28 Barendregt JJ, Doi SA, Lee YY, Norman RE, Vos T. Meta-analysis of prevalence. *J Epidemiol Community Health* 2013; **67**: 974–78.
- 29 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**: 177–88.
- 30 Borenstein M. Introduction to meta-analysis. Chichester: John Wiley & Sons, 2009.
- 31 Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**: 1539–58.
- 32 R core team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing, 2017.
- 33 UN Population Division. World population prospects 2017. 2018. <https://population.un.org/wpp/> (accessed July 10, 2018).
- 34 StataCorp. Stata statistical software: release 14. College Station: StataCorp LP, 2014.
- 35 Kadi Z, Bouguermouh A, Djenaoui T, Allouache A, Dali S, Hadji N. Chlamydial genital infection in Algiers: a sero-epidemiological survey. *Trans R Soc Trop Med Hyg* 1990; **84**: 863–65.
- 36 Abdel Monem AA, Bassyoumi MI, Soliman MY, Ahmed SR, Abdel Hafeez HA. *Chlamydia trachomatis* antigen detection in female infertility. *El-Minia Med Bull* 2005; **16**: 104–12.
- 37 Aboul Atta HNE, Ibrahim AA. Role of chlamydia and mycoplasma in the etiology of non-gonococcal urethritis in men. *Egypt J Med Microbiol* 1995; **4**: 355–60.
- 38 Badary MS. Study of the role of cervical chlamydial infection in unexplained infertility and mucopurulent cervicitis. *Egypt J Med Microbiol* 1996; **5**: 431–38.
- 39 Berry ME, El Shabrawy A. *Chlamydia trachomatis* infection and relation to female infertility. *Egypt J Med Microbiol* 1996; **5**: 297–304.
- 40 Diab KM. Gonococcal and chlamydial antibodies in Egyptian women with ectopic pregnancy. *New Egypt J Med* 1993; **8**: 1006–10.
- 41 Draz EI, Hassan AM, Khalil HS, Elomary MA. Evaluation of pelvic inflammatory disease potential in cholinesterase inhibitor pesticide-exposed females. *Environ Sci Pollut Res Int* 2018; **25**: 30818–25.
- 42 El-Sayed N, Abdallah M, Abdel Mobdy A, et al. Evaluation of selected reproductive health infections in various Egyptian population groups in Greater Cairo. Cairo: Egypt Ministry of Health and Population, 2002.
- 43 Mosbah A, Nabel Y. *Helicobacter pylori*, *Chlamydiae pneumoniae* and *trachomatis* as probable etiological agents of preeclampsia. *J Matern Fetal Neonatal Med* 2016; **29**: 1607–12.
- 44 Mousa A. The association between *Chlamydia trachomatis* and cervical intra-epithelial neoplasia. *Zagazig Med Assoc J* 1990; **3**: 41–46.
- 45 Nada AM, Hassan FM, Al-Azhary NH. Detection of *Chlamydia trachomatis* in patients with unexplained infertility: a case control study. *Egypt J Med Microbiol* 2015; **24**: 35–38.
- 46 Sullam SA, Mahfouz AA, Dabbous NI, el-Barrawy M, el-Said MM. Reproductive tract infections among married women in Upper Egypt. *East Mediterr Health J* 2001; **7**: 139–46.
- 47 Zaki SA. Prevalence of endocervical genital mycoplasmas and *Chlamydia trachomatis* in infertile, abortive and pregnant women in Alexandria. *Bull Alex Fac Med* 1989; **25**: 1031–44.
- 48 Ahmadi A, Khodabandehloo M, Ramazanzadeh R, et al. The relationship between *Chlamydia trachomatis* genital infection and spontaneous abortion. *J Reprod Infertil* 2016; **17**: 110–16.
- 49 Ahmadi MH, Mirsalehian A, Sadighi Gilani MA, Bahador A, Afraz K. Association of asymptomatic *Chlamydia trachomatis* infection with male infertility and the effect of antibiotic therapy in improvement of semen quality in infected infertile men. *Andrologia* 2018; **50**: e12944.
- 50 Ahmadnia E, Kharaghani R, Maleki A, et al. Prevalence and associated factors of genital and sexually transmitted infections in married women of Iran. *Oman Med J* 2016; **31**: 439–45.
- 51 Badami N, Salari MH. Rate of *Chlamydia trachomatis*, *Mycoplasma hominis* and *Ureaplasma urealyticum* in infertile females and control group. *Iran J Public Health* 2001; **30**: 57–60.
- 52 Baghchesaraei H, Amini B, Hossaini M. Prevalence of infection with *Neisseria gonorrhoeae* and *Chlamydia trachomatis* in women visitors of gynecology and obstetrics clinics in Zanjan Province of Iran. *Afr J Microbiol Res* 2011; **5**: 2447–50.
- 53 Bagheri S, Roghanian R, Golbang N, Golbang P, Esfahani MHN. Molecular evidence of *Chlamydia trachomatis* infection and its relation to miscarriage. *Int J Fertil Steril* 2018; **12**: 152–56.
- 54 Behrooz R. Epidemiology of *Chlamydia trachomatis* infections in pregnant women of the hospitals of Tehran University of Medical Sciences. *J Nurs Midwifery Sci* 2001; **1**: 14–20.
- 55 Chamani-Tabriz L, Jedi TM, Zeraati H, et al. A molecular survey of *Chlamydia trachomatis* infection in married women: a cross sectional study on 991 women. *Tehran Univ Med J* 2008; **66**: 498–91.
- 56 Cheraghi M, Rahimi Z, Parsa S. Prevalence of cervical-vaginal infections in the pap-smear samples in Iran. *Glob J Health Sci* 2014; **16**: 201–06.
- 57 Dehghan ML, Aflatoonian A, Talebi AR, Eley A, Pacey AA. Relationship between *Chlamydia trachomatis* and *Mycoplasma genitalium* infection and pregnancy rate and outcome in Iranian infertile couples. *Andrologia* 2017; **49**.
- 58 Goshayeshi L, Vahid Roudsari F, Ghazvini K, Nomani H, Amel Jamehdar S. Pilot prevalence of *Chlamydia trachomatis* by PCR in female infertile referred to study center of infertility in Mashhad. *Iran South Med J* 2015; **18**: 92–99.
- 59 Haghighi Hasanabad M, Bahador A, Mohammadzadeh M, Haghighi F. Prevalence of *Chlamydia trachomatis*, *Neisseria gonorrhoeae* and *Ureaplasma urealyticum* in pregnant women of Sabzevar, Iran. *Sex Transm Infect* 2013; **89** (suppl 1): A233–34.
- 60 Jahromi AS, Farjam MR, Mogharrab F, et al. *Chlamydia trachomatis* in women with full-term deliveries and women with abortion. *Am J Infect Dis* 2010; **6**: 66–69.
- 61 Javanmard D, Behravan M, Ghannadkafi M, Salehabadi A, Ziaee M, Namaei MH. Detection of *Chlamydia trachomatis* in pap smear samples from south Khorasan province of Iran. *Int J Fertil Steril* 2018; **12**: 31–36.
- 62 Joolayi F, Navidifar T, Jaafari RM, Amin M. Comparison of *Chlamydia trachomatis* infection among infertile and fertile women in Ahvaz, Iran: a case-control study. *Int J Reprod Biomed (Yazd)* 2017; **15**: 713–18.
- 63 Kajbaf MJ, Gholamnezhad A. Prevalence of *Chlamydia trachomatis* antigen and antibody in infertile women in Ahwaz. *Iran Biomed J* 1998; **2**: 45–48.
- 64 Kamyabi Z. The survey of anti *Chlamydia* antibody in infertile women undergoing diagnostic laparoscopy in Afzalipour medical centre, Kerman—Iran. *Int J Gynaecol Obstet* 2009; **107**: S218.
- 65 Khezerdoust S, Hagh Elahi F, Roustaei S, Badami N, Naghizadeh MM, Jafarabadi M. *Chlamydia trachomatis* infection in pregnant women. *J Reprod Infertil* 2009; **10**: 121–28.
- 66 Marashi SM, Moulana Z, Imani Fooladi AA, Mashhadi Karim M. Comparison of genital *Chlamydia trachomatis* infection incidence between women with infertility and healthy women in Iran using PCR and immunofluorescence methods. *Jundishapur J Microbiol* 2014; **7**: e9450.
- 67 Farhoudi B, Kamali K, Rajabpoor Z. Situation analysis of sexually transmitted infections in the Islamic Republic of Iran 2008. Tehran: Ministry of Health and Medical Education, 2008.
- 68 Meidani M. Frequency of *Chlamydia trachomatis* infection in asymptomatic males based on PCR in Tehran, Iran. *Int Urogynecol J Pelvic Floor Dysfunct* 2009; **20** (suppl 3): S460–61.
- 69 Mousavi A, Ramezanzadeh R, Farhadifar F, et al. Detection of *Chlamydia trachomatis* in fertile and infertile women in Sanandaj by PCR. *Iran J Public Health* 2014; **43**: 63.
- 70 Pourabbas B, Rezaei Z, Mardaneh J, Shahian M, Alborzi A. Prevalence of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infections among pregnant women and eye colonization of their neonates at birth time, Shiraz, Southern Iran. *BMC Infect Dis* 2018; **18**: 477.
- 71 Rashidi BH, Chamani-Tabriz L, Haghollahi F, et al. Effects of *Chlamydia trachomatis* infection on fertility; a case-control study. *J Reprod Infertil* 2013; **14**: 67–72.
- 72 Rohi E, Ghasemi K, Kahnemoui Agdam F. Incidence of non-gonococcal infection in childbearing and pregnant women in Ardabil. *Int J Mol Clin Microbiol* 2011; **1**: 71–76.
- 73 Rostami MN, Rashidi BH, Aghsaghloo F, Nazari R. Comparison of clinical performance of antigen-based enzyme immunoassay (EIA) and major outer membrane protein (MOMP)-PCR for detection of genital *Chlamydia trachomatis* infection. *Int J Reprod Biomed (Yazd)* 2016; **14**: 411–20.

- 74 Safdari H, Safdari A, Tahaghoghi S, Yari A, Ghazvini K. Prevalence of *Chlamydia trachomatis* among women with genital infection in northeast of Iran in 2013. *Iran J Obstet Gynecol Infertil* 2015; 18: 1–6.
- 75 Sattari M, Ghiami Rad M, Ghasemzadeh A, Mohammadoghli Reihan Z. Frequency of anti-*Chlamydia trachomatis* antibodies in infertile women referred to Tabriz Al-Zahra hospital. *Int J Reprod Biomed (Yazd)* 2017; 15: 17–20.
- 76 Sisakht AJ, Omidifar N, Mohamadkhani N, Karimpoorfard M, Kargar M, Shokripour M. Assessing the presence of *Chlamydia trachomatis* genome in pregnant women with spontaneous abortion using polymerase chain reaction method in Yasuj: first report from southwest of Iran. *J Educ Health Promot* 2017; 6: 45.
- 77 Yeganeh O, Jjedi-Tehrani M, Yaghmaie F, et al. A survey on the prevalence of *Chlamydia trachomatis* and *Mycoplasma genitalium* infections in symptomatic and asymptomatic men referring to urology clinic of Labbafnejad Hospital, Tehran, Iran. *Iran Red Crescent Med J* 2013; 15: 340–44.
- 78 Zahirnia Z, Mansouri S, Saffari F, Mansouri G. Frequency of *Chlamydia trachomatis*, *Mycoplasma genitalium*, and *Ureaplasma urealyticum* isolated from vaginal samples of women in Kerman, Iran. *Arch Clin Infect Dis* 2018; 13: e65334.
- 79 Abdulkhuder NA, Yousif MG, Sadiq AM. Detection of immunological markers for *Chlamydia trachomatis* and *Trichomonas vaginalis* infection in women with obstetric complications in Najaf, Iraq. *J Al-Qadisiyah Pure Sci* 2014; 19: 1–25.
- 80 Abdul-Karim ET, Abdul-Muhyen N, Al-Saadie M. *Chlamydia trachomatis* and rubella antibodies in women with full-term deliveries and women with abortion in Baghdad. *East Mediterr Health J* 2009; 15: 1407–11.
- 81 Abdullah FT. Study the role of *Chlamydia trachomatis* in ectopic pregnancy in Iraqi women. *J Biotechnol Res Cent* 2012; 6: 19–25.
- 82 Ahmed DW. Effects of interleukin-2 (IL-2) and interleukin-6 (IL-6) in recurrent spontaneous abortion (RSA). *Iraqi J Pharm Sci* 2008; 17: 74–79.
- 83 Al-Hamdani SM, Shemran HA, Kadhim HS, Yasser H. Prevalence of anti-*Chlamydia trachomatis* immunoglobulins in pregnant women in Al-Kadhimiya Teaching Hospital. *J Coll Teach* 2010; 1: 59–70.
- 84 Al-Husseinei RK, Kadaem KM, Al-Janabi LA, Fathei MiM. ELFA and IFAT techniques to detect chlamydial infections in Baghdad women and its effect on the immunoglobulins level. *J Facult Med* 2009; 51: 192–97.
- 85 Ali MH, Al-Kazaz AA. Molecular detection of *Chlamydia trachomatis* infection among males with abnormal semen. *Iraqi J Sci* 2018; 59: 2005–11.
- 86 Alkhafaf D. Prevalence of *Chlamydia trachomatis* in women of Al-Diwaniya city. *J Babylon Univ* 2013; 21: 1232–38.
- 87 Hwaid AH, Hasan A-RS, Al-Duliami AA. Seroprevalence of anti-*C trachomatis* IgG and IgM antibodies among pregnant women in Diyala province. *J Facult Med Baghdad* 2013; 55: 348–51.
- 88 Ismail MK, Ali AS. Evaluation of *Chlamydia trachomatis* antibodies in women with infertility. *Al-Mustansiriyah J Sci* 2012; 23: 21–28.
- 89 Mohammed NAM, Salman AH, Hasan FK. *Chlamydia trachomatis* and recurrent spontaneous abortion in Iraqi pregnant women. *Iraqi J Med Sci* 2012; 10: 42–46.
- 90 Mohammed IH, Al-Awadei SJ, Saadedin SMK. Molecular diagnosis of *Chlamydia trachomatis* in infertile Iraqi women using real time-PCR and comparison with other methods. *Iraqi J Sci* 2017; 58: 1437–46.
- 91 Yahya M, Al-Siraj D. Study of genital *Mycoplasma* and *Chlamydia trachomatis* which presence in infertile males. *Al-Taqani* 2009; 22: A23–31.
- 92 Abusarah EA, Awwad ZM, Charvalos E, Shehabi AA. Molecular detection of potential sexually transmitted pathogens in semen and urine specimens of infertile and fertile males. *Diagn Microbiol Infect Dis* 2013; 77: 283–86.
- 93 Al-Ramahi M, Mahafzah A, Saleh S, Fram K. Prevalence of *Chlamydia trachomatis* infection in infertile women at a university hospital in Jordan. *East Mediterr Health J* 2008; 14: 1148–54.
- 94 As'ad A. National AIDS program, final report sexually transmitted infections (STI) prevalence study, Jordan. 2004.
- 95 Awwad ZM, Al-Amarat AA, Shehabi AA. Prevalence of genital chlamydial infection in symptomatic and asymptomatic Jordanian patients. *Int J Infect Dis* 2003; 7: 206–09.
- 96 Mahafzah AM, Al-Ramahi MQ, Asa'd AM, El-Khateeb MS. Prevalence of sexually transmitted infections among sexually active Jordanian females. *Sex Transm Infect* 2008; 35: 607–10.
- 97 Jordan Ministry of Health. Prevalence of reproductive tract infections in women attending selected urban OB/GYN clinics in Jordan. 2004.
- 98 Al-Awadhi R, Al-Shaheen A, Al-Juwaiser A, George SS, Sharma P, Kapila K. Prevalence of infectious organisms observed in cervical smears between 1997–2014 at Mubarak Al-Kabeer Hospital, Kuwait. *Sultan Qaboos Univ Med J* 2018; 18: e324–28.
- 99 Al-Sweih NA, Khan S, Rotimi VO. Prevalence of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* among asymptomatic women attending the Capital Health region clinics in Kuwait. *Sex Transm Infect* 2011; 38: 793–97.
- 100 Al-Sweih NA, Al-Fadli AH, Omu AE, Rotimi VO. Prevalence of *Chlamydia trachomatis*, *Mycoplasma hominis*, *Mycoplasma genitalium*, and *Ureaplasma urealyticum* infections and seminal quality in infertile and fertile men in Kuwait. *J Andrology* 2012; 33: 1323–29.
- 101 Deeb ME, Awwad J, Yeretizian JS, Kaspar HG. Prevalence of reproductive tract infections, genital prolapse, and obesity in a rural community in Lebanon. *Bull World Health Organ* 2003; 81: 639–45.
- 102 Hancali A, Bellaji B, Jennane S, et al. Trend of STIs prevalence among women and men in Morocco between 1999 to 2011. *Sex Transm Infect* 2015; 91: A155.
- 103 Hulstein SH, Matser A, Alberts CJ, et al. Differences in *Chlamydia trachomatis* seroprevalence between ethnic groups cannot be fully explained by socioeconomic status, sexual healthcare seeking behavior or sexual risk behavior: a cross-sectional analysis in the Healthy Life in an Urban Setting (HELIUS) study. *BMC Infect Dis* 2018; 18: 612.
- 104 Morocco Ministry of Health. Etude de prévalence IST chez les femmes consultantes en SM1/PF à la Wilaya de Rabat, Rapport final, Programme National de lutte contre les IST/SIDA. Rabat: Morocco Ministry of Health, 2001.
- 105 Abu-Raddad LJ. Additional data points from the Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project database. Tehran: Infectious Disease Epidemiology Group, 2017.
- 106 Radouani F, Takourt B, Ibrahimy S, Sekkat S, Guinet R, Benslimane A. Contribution of *Chlamydia trachomatis* infection to infertility. *Rev Fr Gynecol Obstet* 1998; 93: 442–46.
- 107 Takourt B, Radouani F, Benckroun A, et al. Seroprevalence of *Chlamydia trachomatis* infection in STD consultants in Morocco. *Bull Soc Pathol Exot* 1995; 88: 194–98.
- 108 Mir AM, Wajid A, Reichenbach L, Khan M. STI prevalence and associated factors among urban men in Pakistan. *Sex Transm Infect* 2009; 85: 199–200.
- 109 Wasti S, Ashfaq MK, Ishaq R, Hamid R. Prevalence of chlamydial infection in females attending antenatal and family planning clinics in Karachi Pakistan. *Aust N Z J Obstet Gynaecol* 1997; 37: 462–65.
- 110 Al-Thani A, Abdul-Rahim H, Alabsi E, et al. Prevalence of *Chlamydia trachomatis* infection in the general population of women in Qatar. *Sex Transm Infect* 2013; 89 (suppl 3): iii57–60.
- 111 Alzahrani AJ, Obeid OE, Hassan MI, Almulhim AA. Screening of pregnant women attending the antenatal care clinic of a tertiary hospital in eastern Saudi Arabia for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infections. *Indian J Sex Transm Dis AIDS* 2010; 31: 81–86.
- 112 Awad NS, Said MM, Mohamed AA, El-Tarras AE. Detection of some sexually transmitted bacterial infection using molecular genetic technique. *World J Med Sci* 2013; 9: 142–46.
- 113 Bashi SA. Chlamydial infection in Riyadh, Saudi Arabia: a sero-epidemiological survey. *Saudi Med J* 1987; 4: 387–90.
- 114 Ghazi HO, Daghestani MH, Mohamed MF. Seropositivity of *Chlamydia trachomatis* among Saudi pregnant women in Makkah. *J Family Community Med* 2006; 13: 61–64.
- 115 Hossain A. Serologic diagnosis of *Chlamydia trachomatis* infections. *Int J Gynaecol Obstet* 1988; 27: 377–80.
- 116 Kamel RM. Screening for *Chlamydia trachomatis* infection among infertile women in Saudi Arabia. *Int J Womens Health* 2013; 5: 277–84.
- 117 Massoud M, Noweir A, Salah M, Saleh WA. Chlamydial infection in Riyadh, Saudi Arabia. *J Egypt Public Health Assoc* 1991; 66: 411–19.
- 118 Ismail SO, Ahmed HJ, Jama MA, et al. Syphilis, gonorrhoea and genital chlamydial infection in a Somali village. *Genitourin Med* 1990; 66: 70–75.



- 119 Nur YA, Groen J, Elmi AM, Ott A, Osterhaus AD. Prevalence of serum antibodies against bloodborne and sexually transmitted agents in selected groups in Somalia. *Epidemiol Infect* 2000; **124**: 137–41.
- 120 WHO. Central South: the 2004 first national second generation HIV/AIDS/STI sentinel surveillance survey. Geneva: World Health Organization, 2005a.
- 121 WHO. The 2004 first national second generation HIV/AIDS/STI sentinel surveillance survey. Geneva: World Health Organization, 2005b.
- 122 Ahmed MA, Hassan NG, Omer ME, Rostami A, Rayis DA, Adam I. *Helicobacter pylori* and *Chlamydia trachomatis* in Sudanese women with preeclampsia. *J Matern Fetal Neonatal Med* 2018; published online Oct 14. DOI:10.1080/14767058.2018.1536738.
- 123 Almroth L, Elmusharaf S, El Hadi N, et al. Primary infertility after genital mutilation in girlhood in Sudan: a case-control study. *Lancet* 2005; **366**: 385–91.
- 124 Ortashi OM, El Khidir I, Herieka E. Prevalence of HIV, syphilis, *Chlamydia trachomatis*, *Neisseria gonorrhoea*, *Trichomonas vaginalis* and candidiasis among pregnant women attending an antenatal clinic in Khartoum, Sudan. *J Obstet Gynaecol* 2004; **24**: 513–15.
- 125 Alkayer SM, Yazji HN, Khamis A. The prevalence of serum IgG antibody to *Chlamydia trachomatis* in infertile women at the university hospital, Syria. *Res J Pharm Technol* 2017; **10**: 1373–75.
- 126 Ghazal-Aswad S, Badrinath P, Osman NA, Abdul-Khalik S, Raasclou T. Is there a correlation between vaginal chlamydia infection and cervical smear abnormalities? A community-based study in the Al-Ain district, United Arab Emirates. *J Obstet Gynaecol Res* 2006; **32**: 63–67.
- 127 Explorable. Non-probability sampling. 2009. <https://explorable.com/non-probability-sampling> (accessed April 10, 2019).
- 128 Glen S. Probability sampling: definition, types, advantages and disadvantages. 2015. <https://www.statisticshowto.datasciencecentral.com/probability-sampling/> (accessed April 10, 2019).
- 129 Johnston LG. Respondent driven sampling (RDS). 2014. <http://www.lisagjohnston.com/respondent-driven-sampling> (accessed April 10, 2019).
- 130 Darougar S, Aramesh B, Gibson JA, Trehan JD, Jones BR. Chlamydial genital infection in prostitutes in Iran. *Br J Vener Dis* 1983; **59**: 53–55.
- 131 Kassaian N, Ataei B, Yaran M, Babak A, Shoaie P, Ataei M. HIV and other sexually transmitted infections in women with illegal social behavior in Isfahan, Iran. *Adv Biomed Res* 2012; **1**: 5.
- 132 Kazerooni PA, Motazedian N, Motamedifar M, et al. The prevalence of human immunodeficiency virus and sexually transmitted infections among female sex workers in Shiraz, south of Iran: by respondent-driven sampling. *Int J STD AIDS* 2014; **25**: 155–61.
- 133 Mirzazadeh ASM, Khajehkazemi R, Hosseini Hooshyar S, et al. HIV and sexually transmitted infections among female sex workers in Iran: findings from the 2010 and 2015 national surveillance surveys. International AIDS Conference; Durban, South Africa; July 18–22, 2016.
- 134 Bellaji B, Hancali A, Jennane S, et al. Prevalence of *Chlamydia trachomatis*, *Neisseria gonorrhoeae* and *Trichomonas vaginalis* in female sex workers in Morocco. *Sex Transm Infect* 2017; **93** (suppl 2): A100.
- 135 Morocco Ministry of Health. Etude de prévalence des IST chez les femmes qui consultent pour pertes vaginales et/ou douleurs du bas ventre. Rabat: Morocco Ministry of Health, 2008.
- 136 Morocco Ministry of Health. HIV integrated behavioral and biological surveillance surveys. Rabat: Morocco Ministry of Health, 2011.
- 137 Morocco Ministry of Health. Integrated behavioral and biological surveillance survey among men who have sex with men (MSM) in Agadir, Casablanca, Marrakech, and Tangier. Rabat: Morocco Ministry of Health, 2015.
- 138 Hawkes S, Collumbien M, Platt L, et al. HIV and other sexually transmitted infections among men, transgenders and women selling sex in two cities in Pakistan: a cross-sectional prevalence survey. *Sex Transm Infect* 2009; **85** (suppl 2): ii8–16.
- 139 Khan MS, Unemo M, Zaman S, Lundborg CS. HIV, STI prevalence and risk behaviours among women selling sex in Lahore, Pakistan. *BMC Infect Dis* 2011; **11**: 119.
- 140 Osama M. Rate of STI among selected MSM and transgenders in 5 cities of Pakistan. *Sex Transm Infect* 2017; **93** (suppl 2): A18.
- 141 Rehan N, Bokhari A, Nizamani NM, et al. National study of reproductive tract infections among high risk groups of Lahore and Karachi. *J Coll Physicians Surg Pak* 2009; **19**: 228–31.
- 142 Znazen A, Frikha-Gargouri O, Berrajah L, et al. Sexually transmitted infections among female sex workers in Tunisia: high prevalence of *Chlamydia trachomatis*. *Sex Transm Infect* 2010; **86**: 500–05.
- 143 Abdel Aleem H, Moubasher AED, Abdel Megeed E. Seroprevalence of *Chlamydia trachomatis* among infertile couples in Assiut. *Ain-Shams Med J* 1996; **47**: 759–67.
- 144 Abdella RM, Abdelmoaty HI, Elsherif RH, et al. Screening for *Chlamydia trachomatis* in Egyptian women with unexplained infertility, comparing real-time PCR techniques to standard serology tests: case control study. *BMC Womens Health* 2015; **15**: 45.
- 145 Azab H, Hassouna AA. Evaluation of clinical findings and chlamydia antibody titre as predictors of tubal factor infertility. *Med J Cairo Univ* 2008; **76**: 171–75.
- 146 Elkayal NM, Mahmoud NF, Abdalla S. Detection of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in Egyptian women suffering from infertility. *Adv Microbiol* 2015; **5**: 769–79.
- 147 El Sayed AF, Al Bakry MM, Zaitoun MM, AbouTalib AF. Comparative study of hysterosalpingography and chlamydia antibody testing in predicting tubal factor infertility. *Ain-Shams Med J* 1997; **48**: 815–25.
- 148 Inhorn MC, Buss KA. Infertility, infection, and iatrogenesis in Egypt: the anthropological epidemiology of blocked tubes. *Med Anthropol* 1993; **15**: 217–44.
- 149 Makled AK, Elkady OS, Swedan KH, Sammour HM, Mohamed EA. Relationship between serum *Chlamydia trachomatis* antibody titer and tubal block in infertile Egyptian women. *Middle East Fertil Soc J* 2013; **18**: 38–41.
- 150 Sadek S, Abo Gabal A, El Hefnawi N. Chlamydial infection and sperm antibodies. *New Egypt J Med* 1993; **8**: 1461–63.
- 151 Siam EM, Hefzy EM. The relationship between antisperm antibodies prevalence and genital *Chlamydia trachomatis* infection in women with unexplained infertility. *Middle East Fertil Soc J* 2012; **17**: 93–100.
- 152 Younis HA, El Sayed SA, Abdel Wahab A, Ayoub AA. Detection of *Chlamydia trachomatis* DNA by polymerase chain reaction in urine of women with tubal infertility as diagnosed by hysterosalpingography and laparoscopy. *Sci J Al-Azhar Med Fac* 2000; **21**: 731–44.
- 153 Zaitoun MM, Zaitoun A. *Chlamydia trachomatis* infection in cases of infertility due to tubal factor: a serological study. *Zagazig Med Assoc J* 1990; **3**: 125–33.
- 154 Zytoon MM. The role of *Chlamydia trachomatis* infection in female infertility. *New Egypt J Med* 1994; **11**: 938–42.
- 155 Golshani M, Eslami G, Ghobadloo SM, et al. Detection of *Chlamydia trachomatis*, *Mycoplasma hominis* and *Ureaplasma urealyticum* by multiplex PCR in semen sample of infertile men. *Iran J Public Health* 2007; **36**: 50–57.
- 156 Hajikhani B, Motallebi T, Norouzi J, et al. Classical and molecular methods for evaluation of *Chlamydia trachomatis* infection in women with tubal factor infertility. *J Reprod Infertil* 2013; **14**: 29–33.
- 157 Kalantar SM, Kazemi MJ, Sheikhha MH, Aflatoonian A, Kafizadeh F. Detection of *Chlamydia trachomatis* infection in female partners of infertile couples. *Iran J Fertil Steril* 2007; **1**: 79–84.
- 158 Mansour Ghanaie M. Relative frequency of *Chlamydia trachomatis* infection in infertile women. *Int J Fertil Steril* 2014; **8**: 110.
- 159 Moazenchi M, Totonchi M, Salman Yazdi R, et al. The impact of *Chlamydia trachomatis* infection on sperm parameters and male fertility: a comprehensive study. *Int J STD AIDS* 2018; **29**: 466–73.
- 160 Nan Bakhsh F, Tarvati MR, Yekta Z, Gol MS, Mehrzad Sadaghiani M, Shokri S. Evaluation of the prognostic value of *Chlamydia trachomatis* antibody testing (CAT), HSG and laparoscopy in predicting tubal factor infertility (TFI) in Motahary Hospital, Urmiah, Iran. *Med J Tabriz Univ Med Sci* 2008; **30**: 119–25.
- 161 Nikbakht R, Saharkhiz N, Ghalambor Dezfouli F. Comparison of level of antibodies against *Chlamydia trachomatis* in infertile women due to tubal factors and fertile women. *J Shahid Sadoughi Univ Med Sci Health Serv* 2008; **16**: 10–15.
- 162 Peivandi S, Moslemizadeh N, Gharajeh S, Ajami A. The role of *Chlamydia trachomatis* IgG antibody testing in predicting tubal factor infertility in northern Iran. *Int J Fertil Steril* 2009; **3**: 143–48.

- 163 Rashidi B, Shahrokh Tehraninezhad E, Alaie F. Prevalence of antichlamydia antibody status in infertile patients. *Int J Fertil Steril* 2007; **1**: 15–18.
- 164 Sadrpour P, Bahador A, Asgari S, Bagheri R, Chamani-Tabriz L. Detection of *Chlamydia trachomatis* and *Mycoplasma genitalium* in semen samples of infertile men using multiplex PCR. *Tehran Univ Med J* 2013; **70**: 623–29.
- 165 Shiahkali AP, Amini K. Identification of herpes simplex virus, *Chlamydia trachomatis*, and *Mycoplasma genitalium* in infertile seminal fluid samples using multiplex-PCR in Kerman province, Iran (2016). *Iran J Obstet Gynecol Infertil* 2018; **21**: 69–74.
- 166 Abid A-AJ, Al-Zwaid AJ. Chlamydial antibodies, pro-inflammatory cytokines and bacterial significance among infertile women. *Med J Babylon* 2015; **12**: 943–47.
- 167 Ahmed ST. Detection of *Chlamydia trachomatis* using polymerase chain reaction (PCR). *Al-Mustansiriyah J Sci* 2012; **23**: 35–42.
- 168 Al-Kattan SAA-K, Mohammed RA. Laparoscopic finding and *Chlamydia trachomatis* infection in infertile women. *Med J Tikrit* 2013; **19**: 315–24.
- 169 Dawood SD. *Chlamydia trachomatis* in Iraqi women with infertility. *Iraqi J Biotechnol* 2011; **10**: 89–94.
- 170 Al Subhi T, Al Jashnmi RN, Al Khaduri M, Gowri V. Prevalence of tubal obstruction in the hysterosalpingogram of women with primary and secondary infertility. *J Reprod Infertil* 2013; **14**: 214–16.
- 171 Qayum M, Khalid-bin-Saleem M. Prevalence of *Chlamydia trachomatis* among asymptomatic women. *J Ayub Med Coll Abbottabad* 2013; **25**: 28–30.
- 172 Al-Hindi A, Al-Helou T, Al-Helou Y. Seroprevalence of *Toxoplasma gondii*, cytomegalovirus, rubella virus and *Chlamydia trachomatis* among infertile women attending in vitro fertilization center, Gaza strip, Palestine. *J Egypt Soc Parasitol* 2010; **40**: 451–58.
- 173 Abdul Jabbar HS. Chlamydial infection in Saudi infertile patients. *Bahrain Med Bull* 1990; **12**: 4–7.
- 174 Alfarrarj DA, Somily AM, Alssum RM, Abotalib ZM, El-Sayed AA, Al-Mandee HH. The prevalence of *Chlamydia trachomatis* infection among Saudi women attending the infertility clinic in Central Saudi Arabia. *Saudi Med J* 2015; **36**: 61–66.
- 175 Sabra SMM, Al-Harbi MS. An influential relationship of seminal fluid microbial infections and infertility, Taif Region, KSA. *World J Med Sci* 2014; **10**: 32–37.
- 176 Gdoura R, Daoudi F, Bouzid F, et al. Detection of *Chlamydia trachomatis* in semen and urethral specimens from male members of infertile couples in Tunisia. *Eur J Contracept Reprod Health Care* 2001; **6**: 14–20.
- 177 Gdoura R, Keskes-Ammar L, Bouzid F, Eb F, Hammami A, Orfila J. *Chlamydia trachomatis* and male infertility in Tunisia. *Eur J Contracept Reprod Health Care* 2001; **6**: 102–07.
- 178 Gdoura R, Kchaou W, Ammar-Keskes L, et al. Assessment of *Chlamydia trachomatis*, *Ureaplasma urealyticum*, *Ureaplasma parvum*, *Mycoplasma hominis*, and *Mycoplasma genitalium* in semen and first void urine specimens of asymptomatic male partners of infertile couples. *J Andrology* 2008; **29**: 198–206.
- 179 Sellami H, Znazen A, Sellami A, et al. Molecular detection of *Chlamydia trachomatis* and other sexually transmitted bacteria in semen of male partners of infertile couples in Tunisia: the effect on semen parameters and spermatozoa apoptosis markers. *PLoS One* 2014; **9**: e98903.
- 180 Ahmadi A, Khodabandehloo M, Ramazanadeh R, et al. The relationship between *Chlamydia trachomatis* genital infection and spontaneous abortion. *J Reprod Infertil* 2016; **17**: 110–16.
- 181 Bagheri S, Roghanian R. Evaluation of *Chlamydia trachomatis* incidence in women who had abortion in Isfahan. *Iran J Public Health* 2014; **43**: 63.
- 182 Massiha A, Khoshkholgh Pahlaviani MRM, Sedighi Kasmaie P, Vand Yousefi J. Study of the reasons of infectious abortions in the subjects who approached Kamali Hospital, Karaj in Iran. *J Reprod Med Endocrinol* 2010; **7**: 346.
- 183 Salari MH, Badami N. The rate of *Chlamydia trachomatis*, *Mycoplasma hominis* and *Ureaplasma urealyticum* in females and habitual abortion. *ACTA Medica Iranica* 2002; **40**: 79–82.
- 184 Al-Nuaimy WA, Al-Jandeel T. Immunological and molecular study of *Chlamydia trachomatis* as causative agent of abortion in Al-Muthanna province. *Iraqi J Vet Sci* 2018; **42**: 99–104.
- 185 Salman YJ. Chlamydia trachomatous antibodies cross reaction with seropositive *Toxoplasma gondii* and cytomegalovirus among women with abortion and outcomes of congenital abnormalities in Kirkuk City. *Tikrit J Pure Sci* 2016; **21**: 1–5.
- 186 Harbord RM, Higgins JPT. Meta-regression in Stata. *Stata J* 2008; **8**: 493–519.
- 187 Dargham SR, Nasrallah GK, Al-Absi ES, et al. Herpes simplex virus type 2 seroprevalence among different national populations of Middle East and north African men. *Sex Transm Infect* 2018; **45**: 482–87.
- 188 Abu-Raddad LJ, Schiffer JT, Ashley R, et al. HSV-2 serology can be predictive of HIV epidemic potential and hidden sexual risk behavior in the Middle East and north Africa. *Epidemics* 2010; **2**: 173–82.
- 189 Vaccarella S, Bruni L, Seoud M. Burden of human papillomavirus infections and related diseases in the extended Middle East and north Africa region. *Vaccine* 2013; **31** (suppl 6): G32–44.
- 190 Rowley J, Hoorn SV, Korenromp E, et al. Chlamydia, gonorrhoea, trichomoniasis and syphilis: global prevalence and incidence estimates, 2016. *Bull World Health Organ* 2019, published online July 4. ID:BLT.18.218651.
- 191 Sonnenberg P, Clifton S, Beddows S, et al. Prevalence, risk factors, and uptake of interventions for sexually transmitted infections in Britain: findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). *Lancet* 2013; **382**: 1795–806.
- 192 CDC, National Center for Health Statistics. National health and nutrition examination survey data, 1999–2014. <https://www.cdc.gov/nchs/nhanes/continuousnhanes/default.aspx> (accessed Sept 25, 2017).
- 193 Redmond SM, Alexander-Kisslig K, Woodhall SC, et al. Genital chlamydia prevalence in Europe and non-European high income countries: systematic review and meta-analysis. *PLoS One* 2015; **10**: e0115753.
- 194 Toomey KE, Rafferty MP, Stamm WE. Unrecognized high prevalence of *Chlamydia trachomatis* cervical infection in an isolated Alaskan Eskimo population. *JAMA* 1987; **258**: 53–56.
- 195 Walsh MS, Hope E, Isaia L, et al. Prevalence of *Chlamydia trachomatis* infection in Samoan women aged 18 to 29 and assessment of possible risk factors: a community-based study. *Trans R Soc Trop Med Hyg* 2015; **109**: 245–51.
- 196 Stamm WE. *Chlamydia trachomatis* infections of the adult. In: Holmes KK, Sparling FP, Stamm WE, et al, eds. Sexually transmitted diseases, 4th edn. New York: McGraw Hill Medical, 2008: 575–93.
- 197 Molano M, Meijer CJ, Weidnerpass E, et al. The natural course of *Chlamydia trachomatis* infection in asymptomatic Colombian women: a 5-year follow-up study. *J Infect Dis* 2005; **191**: 907–16.
- 198 Heijne JC, Herzog SA, Althaus CL, Low N, Kretzschmar M. Case and partnership reproduction numbers for a curable sexually transmitted infection. *J Theor Biol* 2013; **331**: 38–47.
- 199 Omori R, Chemaitelly H, Althaus CL, Abu-Raddad LJ. Does infection with *Chlamydia trachomatis* induce long-lasting partial immunity? Insights from mathematical modelling. *Sex Transm Infect* 2019; **95**: 115–21.
- 200 Al-Mutairi N, Joshi A, Nour-Eldin O, Sharma AK, El-Adawy I, Rijhwani M. Clinical patterns of sexually transmitted diseases, associated sociodemographic characteristics, and sexual practices in the Farwaniya region of Kuwait. *Int J Dermatol* 2007; **46**: 594–99.
- 201 Ismail SO, Ahmed HJ, Grillner L, Hederstedt Issa BA, Bygdeman S. Sexually transmitted diseases in men in Mogadishu, Somalia. *Int J STD AIDS* 1990; **1**: 102–06.
- 202 Faisal A, Cleland J. Migrant men: a priority for HIV control in Pakistan? *Sex Transm Infect* 2006; **82**: 307–10.
- 203 Mumtaz GR, Riedner G, Abu-Raddad LJ. The emerging face of the HIV epidemic in the Middle East and north Africa. *Curr Opin HIV AIDS* 2014; **9**: 183–91.
- 204 Inhorn MC. Quest for conception: gender, infertility and Egyptian medical traditions. Philadelphia, PA: University of Pennsylvania Press, 1994.
- 205 Zurayk H, Sholkamy H, Younis N, Khattab H. Women's health problems in the Arab world: a holistic policy perspective. *Int J Gynaecol Obstet* 1997; **58**: 13–21.
- 206 Eggert-Kruse W, Buhlinger-Gopfarth N, Rohr G, et al. Antibodies to *Chlamydia trachomatis* in semen and relationship with parameters of male fertility. *Hum Reprod* 1996; **11**: 1408–17.

- 207 Eggert-Kruse W, Weltin M, Strowitzki T. Are chlamydial lipopolysaccharide-directed antibodies in seminal plasma or serum clinically significant during investigation of male infertility? *Urology* 2011; **77**: 1101–06.
- 208 Dietrich W, Rath M, Stanek G, Apfalter P, Huber JC, Tempfer C. Multiple site sampling does not increase the sensitivity of *Chlamydia trachomatis* detection in infertility patients. *Fertil Steril* 2010; **93**: 68–71.
- 209 Miron N, Socolov D, Mares M, et al. Bacteriological agents which play a role in the development of infertility. *Acta Microbiol Immunol Hung* 2013; **60**: 41–53.
- 210 Faour M. Fertility policy and family planning in the Arab countries. *Stud Fam Plann* 1989; **20**: 254–63.
- 211 Rashad H, Khadr Z. The demography of the Arab region: new challenges and opportunities. In: Human capital: population economics in the Middle East. Sirageldin I, ed. London: I B Tauris, 2002: 37–49.
- 212 Rashad H, Osman M. Nuptiality in Arab countries: changes and implications. In: Cairo papers in social science. Hopkins NS, ed. Cairo: American University in Cairo Press, 2003; 20–50.
- 213 Bristow CC, Mathelier P, Ocheretina O, et al. *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, and *Trichomonas vaginalis* screening and treatment of pregnant women in Port-au-Prince, Haiti. *Int J STD AIDS* 2017; **28**: 1130–34.
- 214 Cabeza J, Garcia PJ, Segura E, et al. Feasibility of *Chlamydia trachomatis* screening and treatment in pregnant women in Lima, Peru: a prospective study in two large urban hospitals. *Sex Transm Infect* 2015; **91**: 7–10.
- 215 Joseph Davey DL, Shull HI, Billings JD, Wang D, Adachi K, Klausner JD. Prevalence of curable sexually transmitted infections in pregnant women in low- and middle-income countries from 2010 to 2015: a systematic review. *Sex Transm Infect* 2016; **43**: 450–58.
- 216 Nguyen M, Le GM, Nguyen HTT, Nguyen HD, Klausner JD. Acceptability and feasibility of sexually transmissible infection screening among pregnant women in Hanoi, Vietnam. *Sex Health* 2019; **16**: 133–38.
- 217 Wynn A, Ramogola-Masire D, Gaolebale P, et al. Prevalence and treatment outcomes of routine *Chlamydia trachomatis*, *Neisseria gonorrhoeae* and *Trichomonas vaginalis* testing during antenatal care, Gaborone, Botswana. *Sex Transm Infect* 2018; **94**: 230–35.
- 218 Syred J, Engler B, Campbell L, Baraitser P, Sheringham J. Exploration of gender differences of *Chlamydia trachomatis* infection amongst young people reveals limitations of using sexual histories to assess risk in high-prevalence areas. *Int J STD AIDS* 2014; **25**: 564–70.
- 219 Horner PJ, Wills GS, Reynolds R, et al. Effect of time since exposure to *Chlamydia trachomatis* on chlamydia antibody detection in women: a cross-sectional study. *Sex Transm Infect* 2013; **89**: 398–403.
- 220 Hocking JS, Temple-Smith M, Guy R, et al. Population effectiveness of opportunistic chlamydia testing in primary care in Australia: a cluster-randomised controlled trial. *Lancet* 2018; **392**: 1413–22.
- 221 WHO. Strategies and laboratory methods for strengthening surveillance of sexually transmitted infections 2012. Geneva: World Health Organization, 2012. <http://www.who.int/reproductivehealth/publications/rtis/9789241504478/en/> (accessed Aug 9, 2014).
- 222 Bozicevic I, Riedner G, Calleja JM. HIV surveillance in MENA: recent developments and results. *Sex Transm Infect* 2013; **89** (suppl 3): iii11–16.
- 223 Ayoub HH, Awad SF, Abu-Raddad LJ. Use of routine HIV testing data for early detection of emerging HIV epidemics in high-risk subpopulations: a concept demonstration study. *Infect Dis Model* 2018; **3**: 373–84.
- 224 Reintjes R, Wiessing L. 2nd-generation HIV surveillance and injecting drug use: uncovering the epidemiological ice-berg. *Int J Public Health* 2007; **52**: 166–72.
- 225 Hessini L. Abortion and Islam: policies and practice in the Middle East and north Africa. *Reprod Health Matters* 2007; **15**: 75–84.
- 226 Shapiro GK. Abortion law in Muslim-majority countries: an overview of the Islamic discourse with policy implications. *Health Policy Plan* 2014; **29**: 483–94.